

# Effect of dietary protein level on the reproductive performance of female of Green catfish ( Hemibagrus nemurus Bagridae )

*by* Indra Suharman

---

**Submission date:** 08-Apr-2023 03:12PM (UTC+0700)

**Submission ID:** 2058930591

**File name:** Journal\_of\_ARD\_2015.pdf (1.66M)

**Word count:** 4613

**Character count:** 23742



## Effect of Dietary Protein Level on the Reproductive Performance of Female of Green Catfish (*Hemibagrus nemurus* Bagridae )

Netti Aryani\* and Indra Suharman

Department of Aquaculture, Fisheries and Marine Science Faculty of Riau University, Panam, Pekanbaru Riau, Indonesia

### Abstract

A study to determine the effect of increasing levels of dietary protein on green catfish (*Hemibagrus nemurus*), was conducted. Four isocaloric semi-purified diets containing 20%, 27%, 32%, and 37% dietary protein were tested. Broodstock performance was evaluated based on growth parameters, reproductive parameters, and larvae production of female broodstock. Results showed that while the 20%, 27%, 32% and 37% protein produced the lowest specific growth rate (SGR) values, there was significant difference between 20%, 27%, 32% and 37% dietary treatments. The 20% dietary treatment also displayed longest time matured gonads and lowest of somatic ovi index and relative fecundity. Larvae production was highest from females fed with 32% and 37% protein, followed by the 27% protein while the diet containing 20% protein produced lowest number of larvae. Based on our results, we suggest that a minimum of 32% protein be included in the diet of female green catfish broodstock.

**Keywords:** *Hemibagrus nemurus*; Broodstock nutrition; Growth; Fecundity; Larvae

### Introduction

Indonesia is one of the countries that are rich in biodiversity and recognized internationally as a 'hot megadiversity country'. Within 1,200 species of freshwater [1,2]. There are 260 species in inland waters include rivers, oxbow lake, lakes, reservoirs and floodplain in Riau Province [3]. In general, the species diversity has been decreased, Siak river was only found 36 species [4], Kampar Kiri river 86 species [5], Oxbow lake Serai 20 species [6], Kampar Kanan river 58 species [7], Kampar Kanan River 36 species [8], Koto Panjang reservoir 26 species [9] and small Giam Siak 37 species [10].

One species of fish that live in the inland waters of Riau Province which has important economic value is green catfish. This fish is highly favored by consumers because the thick meat, little prickly and the delicious taste, weight can reach sizes from 750 to 1000 g/individual so as to have a high market value i.e. 60,000 to 75,000 IRD/kg [11]. At the present time in Riau Province has been cultured with intensively by in cages farmers in river basin and ponds. But the rapidly development of cultured green catfish have not balanced by the high production because not supported by larvae production with good quality and quantity [12]. This is due, among others, the difficulty of getting broodstock mature gonads. Moreover, the results showed that green catfish fed with enriched vitamin E ranged from 150 to 450 mg/kg feed resulted in fecundity ranged from 23,750 to 30,000 eggs/kg body weight [13], implantation with LHRH-a is still a little ranged from 20,815 to 32,000 eggs/spawn [13], with 17- $\beta$ -estradiol implantation ranged from 40,875 to 63,724 eggs/spawn [12].

Demand of green catfish juvenile age 50 days for cultured in Pekanbaru City and Kampar Regency is currently as 1,000,000 individuals every year, the majority of juvenile derived from wild fish catch and highly dependent on season and availability stocks. In order available at any time juveniles in quantity and quality for cultured to be supplied from hatcheries [12].

In the research, the author wants to analyze the effectiveness of the dietary protein levels on reproductive performance green catfish, because the nutrition of broodstock is one of the least studied areas because the biological mechanisms, such as gonadal maturation, are extremely complex processes [12,14-19]. The gonadal development

and fecundity are affected by several nutrients, especially in continuous spawning fish with short vitellogenic periods as *Ictalurus punctatus* [18]. In general, the nutritional status of the female can influence gonadal development and limit the amount and the quality of the eggs [15,20,21]. The current study was designed to investigate the effects of dietary protein level on various reproductive aspects of green catfish utilizing semi-purified diets.

### Methodology

#### Broodstock and cages

Adult female and male green catfish (*H. nemurus*) were obtained from a commercial fish farm in Sungai Paku Village, Kampar Regency Riau Province and have been kept for >1.5 years in pond, average weight is 750  $\pm$  100 g/individuals (Figure 1). The female of green catfish to be treated as individuals first twelve adapted for three months in the cages (2  $\times$  1  $\times$  1 m) with a stocking density of individual/cage. All cages are placed in pond (22  $\times$  8  $\times$  2 m) with an average water height of one meter. The pond water temperature ranged from 26°C to 28°C. Feeding was done twice daily and fish were fed a predetermined ration of 5% body weight day<sup>-1</sup>. During the adaptation of fish fed with commercial (pellets) with proximate composition are water content (% dry weight) 12.0%, crude protein 28.0%, lipid 5.5%, crude fiber 6.2% and crude ash 13.0%. Experiments conducted on green catfish female gonad maturity stage one with four groups and three replications. Each group given in the form of pellet feed with protein content of 20%, 27%, 32% and 37% respectively.

\*Corresponding author: Netti Aryani, Department of Aquaculture, Fisheries and Marine Science Faculty of Riau University, Pekanbaru Riau, Indonesia, Tel: 265 888 99 846; E-mail: [nettiaryani@yahoo.com](mailto:nettiaryani@yahoo.com)

Received July 09, 2015; Accepted July 27, 2015; Published August 27, 2015

Citation: Aryani N, Suharman I (2015) Effect of Dietary Protein Level on the Reproductive Performance of Female of Green Catfish (*Hemibagrus nemurus* Bagridae). J Aquac Res Development 6: 377. doi:10.4172/2155-9546.1000377

Copyright: © 2015 Aryani N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

The spawning green catfish conducted with GnRH $\alpha$  stimulation with dopamine antagonist at a dose of 0.5 ml/kg body weight and conducted artificial spawning. Hatching eggs each treatment and replication carried on fiber tubs (200 × 80 × 40 cm), height of water in the fiber tubs is 20 cm, water temperature during embryonic development ranged from 28 to 29°C, pH 6.8 to 7 and dissolved oxygen 5.6 to 6.8 mg/l. Egg samples from each treatment and replications preserved with Gilson solution. Gilson is preserved with a solution consisting 100 ml of 60% alcohol, 880 ml of distilled water, 15 ml of nitric acid, 18 ml lacial acetic acid and 20 grams of mercury chloride. Furthermore, the diameter of the eggs was measured with a microscope Olympus CX 21 to 30 eggs of from each treatment and replications. The water temperature, dissolved oxygen (DO), and pH levels of the test chambers were regularly monitored.

### Diets

Four semi-purified isocaloric diets with an increasing level of dietary protein (20%, 27%, 32%, and 37%) were formulated utilizing fish meal and soybean meal as protein sources. Proximate analysis of diets was conducted according to Saraswanti Indo Genetech (2013) for verification of nutrient levels.

### Checking the oocytes maturation

All fish were individually marked using floy-tags and weighed. Oocytes sampled in vivo were taken from females using the method described by Syandri [22], and were placed in Serra's solution (6:3:1, 70% ethanol, 40% formaldehyde and 99.5% acetic acid) for clarification of the cytoplasm. After 5 min, the position of oocytes nucleus was determined using a four-stage scale:

stage 1 germinal vesicle in central position

stage 2 early migration of germinal vesicle (less than half of radius)

| Ingredient                      | Dietary protein levels (%) |       |      |      |
|---------------------------------|----------------------------|-------|------|------|
|                                 | 20%                        | 27%   | 32%  | 37%  |
| Fish meal (60% CP) <sup>a</sup> | 21                         | 30    | 36   | 40   |
| soybean meal                    | 10                         | 12    | 15   | 18   |
| fine bran                       | 20                         | 20    | 20   | 20   |
| Def 24                          | 25                         | 20    | 15   | 5    |
| Shark liver oil                 | 5                          | 5     | 5    | 5    |
| Corn oil                        | 3                          | 3     | 3    | 3    |
| Vitamin premix <sup>b</sup>     | 2.5                        | 1.5   | 1.5  | 1.5  |
| Mineral premix <sup>c</sup>     | 2.5                        | 1.5   | 1.5  | 1.5  |
| CMC                             | 1.5                        | 1.5   | 1.5  | 1.5  |
| Cellulose                       | 8.5                        | 5.5   | 5    | 4.5  |
| <i>Proximat (dry weight)</i>    |                            |       |      |      |
| Crude protein (%)               | 20.2                       | 26.05 | 31.8 | 37.1 |
| Crude lipid (%)                 | 9.5                        | 9.8   | 10.1 | 10.3 |
| Ash (%)                         | 5.60                       | 5.62  | 5.64 | 5.67 |
| crude fiber (%)                 | 4.5                        | 4.6   | 4.70 | 4.75 |
| Energy (k cal/g)                | 4.60                       | 4.65  | 4.65 | 4.66 |

<sup>a</sup>Anugrah Sakti Fish Meal (crude protein: 60.0%, crude lipid: 13.0%, crude fiber: 1.5%, crude ash: 25%).

<sup>b</sup>Vitamin mix (mg/100 g feed): Thiamin-HCl 5.0; riboflavin 5.0; Ca-pantothenate 10.0; niacin 2.0; pyridoxin-HCl 4.0; biotin 0.6; folic acid 1.5; cyanocobalamin 0.01; inositol 200; p-aminobenzoic acid 5.0; menadion 4.0; vit A palmitat 15.0; cholecalciferol 1.9;  $\alpha$ -tocopherol 20.0; cholin chloride 900.0

<sup>c</sup>Mineral mix (mg/100 g feed): KH<sub>2</sub>PO<sub>4</sub> 412; CaCO<sub>3</sub> 282; Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub> 618; FeCl<sub>3</sub>·4H<sub>2</sub>O 166; ZnSO<sub>4</sub>·9.99; MnSO<sub>4</sub>·6.3; CuSO<sub>4</sub>·2; CoSO<sub>4</sub>·7H<sub>2</sub>O 0.05; KJ 0.15; Dekstrin 450; Selulosa 53.51.

Table 1: Composition and proximate analysis respectively of test diets.

stage 3 late migration of germinal vesicle (more than half of radius)

stage 4 periphery germinal vesicle or germinal vesicle breakdown, GVBD (Table 1).

### Parameters analyzed were

Specific growth rate (SGR %/day):  $[(\ln W_t - \ln W_i) / T] \times 100$  where  $W_t$ =mean final weight,  $W_i$ =mean initial weight and  $T$ =total experimental days. Feed conversion ratio (FCR): total feed fed (g)/total wet weight gain (g). Time mature gonadal calculated from the time the fish began dietary protein levels granted until the fish reaches a mature gonadal [days]. Ovi somatic index (OSI) was determined using the expression:  $OSI = BTO / BW \times 100\%$ ; where BTO: ovulation egg weight and BW: Body Weight. Absolute fecundity (AF)=OVA × GW; where OVA: oocyte number per ovary gram and GW: Gonadal Weight. Relative fecundity was estimated using the formula:  $RF = AF / BW$ ; where AF: Absolute Fecundity; BW: Body Weight.

### Chemical analysis

The proximate composition of egg samples from each treatment analysed using the following methods: moisture content by drying in an oven at 100°C for 24 h; protein using a Kjeltec Auto 1030 (Tecator Ltd.); and lipid by the Soxhlet method (AOAC., 1995). Egg samples for amino acid analyses were stored at -70°C before use. Total fry production: Total larvae harvested throughout experimental period (age 14 days).

### Statistical analyses

Data were analysed by one-way analysis of variance (ANOVA) and then tested using Dunnett-s test. All statistical analyses were performed using SPSS 17. (SPSS, Tokyo, Japan). In all tests, the level of significance used was 0.05.

## Results and Discussion

### Final weight, SGR and FCR

Growth of green catfish females dietary protein levels showed that while there was significantly ( $p < 0.05$ ) difference between final weight of diets 20%, 27%, 32% and 37% (Table 2 and Figure 1). Weight gain was lowest for the 20% protein diet, followed by 27%, 32% and 37% protein diets. Specific growth rate (SGR) was also significantly different between protein diet, SGR was lowest for the 20% protein diet, followed by the



Figure 1: Broodstock of *H.nemurus*.

27%, 32% and 37% dietary protein levels. Results from the FCR values also showed that the 32% and 37% dietary protein levels produced the highest efficiency in feed utilization (Table 2) (Figures 1-4).

### The time sexual maturity and percentage of eggs weight

The mature gonadal of green catfish females dietary protein levels showed that while there was significantly ( $p < 0.05$ ) difference between the time sexual maturity (Figure 2 and Table 3). Average egg ovulation (SOI) was also significantly different between protein diet, SOI was lowest for the 20%, followed by the 27%, 32% and 37% dietary protein levels (Table 3).

### Fecundity and eggs diameter

The mean absolute fecundity of green catfish females ranged from 40,033 to 64,650 eggs/spaw (Table 4). Absolute fecundity tended to increase with increasing level of diet protein. Increasing dietary protein levels also increase the relative fecundity and eggs diameter. The absolute fecundity, relative fecundity and eggs diameter were significantly ( $p < 0.05$ ) among the dietary protein levels (Table 4).

|                    | Dietary protein levels      |                            |                             |                            |
|--------------------|-----------------------------|----------------------------|-----------------------------|----------------------------|
|                    | 20%                         | 27%                        | 32%                         | 37%                        |
| Initial weight (g) | 754.33 ± 14.50              | 755 ± 5                    | 747 ± 11.53                 | 754.66 ± 5.03              |
| Final weight (g)   | 799.82 ± 13.50 <sup>a</sup> | 813.14 ± 6.89 <sup>b</sup> | 817.85 ± 13.56 <sup>c</sup> | 841.45 ± 6.82 <sup>d</sup> |
| Weight gain (g)    | 44.82 ± 0.36 <sup>a</sup>   | 58.14 ± 2.35 <sup>b</sup>  | 70.85 ± 2.68 <sup>c</sup>   | 86.79 ± 2.54 <sup>d</sup>  |
| SGR (%)            | 0.1 ± 0.002 <sup>a</sup>    | 0.16 ± 0.008 <sup>b</sup>  | 0.28 ± 0.02 <sup>c</sup>    | 0.41 ± 0.03 <sup>d</sup>   |
| FCR                | 2.74 ± 0.16 <sup>a</sup>    | 2.05 ± 0.14 <sup>b</sup>   | 1.36 ± 0.18 <sup>c</sup>    | 1.18 ± 0.07 <sup>d</sup>   |

Table 2: Effect of different dietary protein levels on various growth parameters of female green catfish (mean ± SE).

| Dietary protein levels | Average body weight (g) | Fecundity (number of eggs per spaw) | Relative fecundity (number of eggs per g gonadal weight) | Eggs diameter (mm)       |
|------------------------|-------------------------|-------------------------------------|--|--------------------------|
| 20%                    | 799.82 ± 13.50          | 40,033 ± 1,789 <sup>a</sup>         | 49 ± 2 <sup>a</sup>                                      | 1.07 ± 0.01 <sup>a</sup> |
| 27%                    | 813.14 ± 6.89           | 46,896 ± 1,300 <sup>b</sup>         | 57 ± 2 <sup>b</sup>                                      | 1.11 ± 0.02 <sup>b</sup> |
| 32%                    | 817.85 ± 13.56          | 58,106 ± 2,393 <sup>c</sup>         | 70 ± 3 <sup>c</sup>                                      | 1.16 ± 0.01 <sup>c</sup> |
| 37%                    | 841.45 ± 6.82           | 64,650 ± 2,107 <sup>d</sup>         | 77 ± 2 <sup>d</sup>                                      | 1.21 ± 0.02 <sup>d</sup> |

Table 3: Effect dietary protein levels on time matured the gonads and percentage of eggs weight (mean ± SE).



Figure 2: Ovary stage IV of *H. nemurus*.



Figure 3: Larvae fourteen days.



Figure 4: Juvenile age 50 days.

### Chemical composition of eggs and total larvae production

Percent protein, lipid and water content in the largest group of oocytes of stage 4 ovaries (post-vitellogenic oocytes; diameter ranged from 1.07 to 1.21 mm, Table 5). Protein and lipid (by dry weight) and water content ranged from 55.76 to 61.21% and 4.71 to 7.33%, respectively, and there was significantly differences ( $P < 0.05$ ) among the different dietary protein levels.

Mean total larvae production increased with an increasing dietary protein level. The highest larvae production was obtained with dietary protein levels of 32% (40,093 ± 2,100 individuals) and 37% (45,255 ± 2,950 individuals), followed by 27% (31,185 ± 3,105 individuals) and 20% (25,621 ± 2,659 individuals). The total larvae production were significantly ( $p < 0.05$ ) among the dietary protein levels (Figures 3 and 4 and Table 5).

### Discussion

The green catfish (*H. nemurus*) females given dietary protein levels 20%, 27%, 32% and 37% can increase weight gain, specific growth rate and decrease feed conversion ratio. In contrast to reddaw crayfish (*Cherax quadricarinatus*) with dietary protein levels 22%, 27%, 32%, and 37% did not affect the final weight [17]. Numerous studies elsewhere have shown that an important contribution of dietary protein toward broodstock performance is in the effect on the body size. Fish

| Dietary protein levels | Average body weight (g) | Fecundity (number of eggs per spawn) | Relative fecundity (number of eggs per g gonadal weight) | Eggs diameter (mm)       |
|------------------------|-------------------------|--------------------------------------|--|--------------------------|
| 20%                    | 799.82 ± 13.50          | 40.033 ± 1.789 <sup>a</sup>          | 49 ± 2 <sup>a</sup>                                      | 1.07 ± 0.01 <sup>a</sup> |
| 27%                    | 813.14 ± 6.89           | 46.896 ± 1.300 <sup>b</sup>          | 57 ± 2 <sup>b</sup>                                      | 1.11 ± 0.02 <sup>b</sup> |
| 32%                    | 817.85 ± 13.56          | 58.106 ± 2.393 <sup>c</sup>          | 70 ± 3 <sup>c</sup>                                      | 1.16 ± 0.01 <sup>c</sup> |
| 37%                    | 841.45 ± 6.82           | 64.650 ± 2.107 <sup>d</sup>          | 77 ± 2 <sup>d</sup>                                      | 1.21 ± 0.02 <sup>d</sup> |

Values with the different superscript in each column are significantly different from each other ( $p < 0.05$ ).

**Table 4:** Effect of different dietary protein levels on fecundity and eggs diameter (mean ± SE).

| Dietary protein levels | Egg protein (%) (dry weight) | Egg lipid (%) (dry weight) | Water content (%) (dry weight) | Average total larvae production |
|------------------------|------------------------------|----------------------------|--------------------------------|---------------------------------|
| 20%                    | 55.76 ± 0.45 <sup>a</sup>    | 21.4 ± 0.55 <sup>a</sup>   | 7.33 ± 0.15 <sup>a</sup>       | 25,621 ± 2,659 <sup>a</sup>     |
| 27%                    | 57.56 ± 0.20 <sup>a</sup>    | 18.33 ± 0.21 <sup>a</sup>  | 6.38 ± 0.10 <sup>a</sup>       | 31,185 ± 3,105 <sup>b</sup>     |
| 32%                    | 58.20 ± 0.4 <sup>a</sup>     | 18.86 ± 0.25 <sup>a</sup>  | 7.27 ± 0.26 <sup>a</sup>       | 40,093 ± 2,100 <sup>c</sup>     |
| 37%                    | 58.29 ± 0.36 <sup>a</sup>    | 17.33 ± 0.73 <sup>a</sup>  | 6.71 ± 0.11 <sup>a</sup>       | 45,255 ± 2,950 <sup>d</sup>     |

Values with the different superscript in each column are significantly different from each other ( $p < 0.05$ ).

**Table 5:** Effect dietary protein levels on percentage protein, lipid (by dry weight) and water content in post-vitellogenic oocytes (diameter 1.9-2.5 mm) of stage 4 ovaries and total larvae production (mean ± SE).

female swordtails (*Xiphophorus helleri*, Poeciliidae) feed with a protein content of 20%, 30%, 40%, 50% and 60% of dietary protein can also enhance final weight and weight gain [15]. In tilapia, female total weight at first maturation increases linearly with higher dietary protein levels [23]. The female tilapia attained puberty and oocyte maturation earlier when feed higher levels of dietary protein and concluded that this was due to the effect of diet on fish growth [20].

The green catfish (*H.nemurus*) females group dietary protein levels exposed can reaches the gonadal mature more rapidly, increased of the index ovi somatic, absolute fecundity, relative fecundity and eggs diameter. With a 37% protein dietary level, capable of reaching a matured gonads of green catfish during 26 days and increase of fecundity as much as 64,650 eggs/individual. While the protein content of feed 20% of the time matured gonads achievement during 58 days and fecundity as much as 40,033 eggs/individual. The female dwarf gourami (*Colisa lalia*) broodstock showed higher total weight and higher number of oocytes undergoing vitellogenesis when feed higher dietary protein level [24]. The lower protein content obtained in both muscle and ovary of female swordtail fed with the 20% protein level probably indicates limited or insufficient protein for maintenance and oocyte development. Continuous feeding with a low dietary protein level also caused female tilapia to utilize body reserves during subsequent spawning seasons compromising muscle deposition and growth [20].

In general the exposed to 20% to 37% of dietary protein levels can increased ovi somatic index of green catfish ranged between 4.47 % to 8.24%. According to [13] that of green catfish exposed LHRHa to 400 µg/kg body weight to produced somatic ovi index 8.08%, and the exposed to 200 to 600 µg/kg body weight of 17β-estradiol can increased ovi somatic index of green catfish ranged between 5.01% to 10.32% [12]. Green catfish were implanted 17β-estradiol at a dose of 200 µg/kg body weight reached matured gonads during 35 days and fecundity as much as 40,875 eggs/individual, whereas at a dose of 600 µg/kg body weight reached matured gonads during 26 days with fecundity

as 52,500 eggs/individual [12]. The *Rhamdia quelen* females were given feed with protein levels of 28%, 34% and 40% had no effect on the main physiological and reproductive parameters. The dietary protein levels 28%, 34% and 40%, produced eggs are 127.00 ± 17.60; 140.00 ± 11.10; 143.00 ± 8.00 egg number ml<sup>-1</sup> spawn respectively. In conclusion, the 28% crude protein dietary level was sufficient for maintenance of the broodstock and the reproductive indexes [19]. The fish reproduction is influenced by the quality of feed among others lipid, protein, fatty acids, vitamins E and C, and carotenoids as major nutrients influencing various reproduction processes such as frequency of spawning and the seed output, fecundity, fertilization, hatching and larval development [14,25], hormone levels 17-β-estradiol [26-28], hormone level IIRH [29,30] and environmental factors [31]. In the present study, percent protein and lipid in the post-vitellogenic oocytes did not vary with the dietary protein level. Similar results were reported by [20] for tilapia (*Oreochromis niloticus*). But the levels of dietary protein significantly affect to larvae production until stage of 14 days. In summary, dietary protein levels had influence on weight gain, specific growth rate, feed conversion ratio, time matured gonadal, index ovi somatic, fecundity, eggs diameter and larvae production.

#### Acknowledgment

Thanks to the Directorate of Research and Community Service (DP2M) Directorate General of Higher Education Ministry of Education which has provided research funding through the National Strategy Research in 2013 and they wish to thank the reviewers at the Journal of Aquaculture Research and Development valuable criticism of the manuscript.

#### References

- Kottelat MAJ, Whitten SN, Kartikasari S, Wirjoatmodjo (1993) Freshwater fishes of western indonesia and sulawesi. Periplus Edition (HK), Jakarta.
- Ministry of Maritime Affairs and Fisheries of the Republic of Indonesia (2012) Rare fish in Indonesia. Directorate General of Marine, Coastal and Small Islands. Ministry of Maritime Affairs and fisheries RI 350.
- Riau Provincial Fisheries Office (1995) Inventory and identification of local fish species specific Riau Province. Fisheries Agency of Riau Province in Figures.
- Iskandar J, Dahiyat Y (2012) The diversity of fish in the Siak River Riau. Bionatura-Journal of Biological Sciences and Physical 1: 51-58.
- Simanjuntak CPH, Rahardjo MF, Sukimin S (2006) Ichthyofauna in Floodplain of Kampar Kiri River. Journal of Ichthyology Indonesia 6: 99-109.
- Alawi H, Rengi T, Tang UM (2008) Directory of commercial fish in inland waters. Bengkalis Regency Riau Province, Unri Press p: 80.
- Fithra RY, Siregar YI (2010) Fishes biodiversity of Kampar River an inventory from Kampar Kanan River. Journal of Environmental Science 2: 139-147.
- Aryani N (2015) Native species in Kampar Kanan River Riau Province Indonesia. International Journal of Fisheries and Aquatic Studies 2: 213-217.
- Warsa A, Nastiti AS, Krismono, Nurfiarini A (2009) Fisheries resources in Koto Panjang Reservoir. Bawal 2: 93-97.
- Aryani N (2001) The use of vitamin E in feed for fish gonad maturation green catfish (*Mystus nemurus*). Journal of Fisheries and Marine Sciences 6: 28-36.
- Marini M, Husnah (2011) The diversity of fish species Giam Siak Kecil Riau Province. In: Proceedings of the Indonesian Open Water Forum 8th. Inland Water Fisheries Research Institute Palembang pp: 396-403.
- Aryani N, Nuraini, Suharman I (2013) Morphological characterization of green catfish (*Hemibagrus nemurus*) aquatic habitat on the different method based truss morfometrics. Journal of Fisheries and Aquaculture 4: 139-142.
- Aryani N, Suharman I (2014) Effects of 17β-estradiol on the reproduction of green catfish (*Hemibagrus nemurus*, Bagridae). Journal of Fisheries and Aquaculture 5: 163-168.
- Aryani N, Syawal H, Bukhari D (2002) The use of LHRH-a for gonadal

- maturation of green catfish (*Mystus nemurus*). Torani 12: 163-168.
15. Izquierdo S, Fernandez-Palacios H, Tacon AGJ (2001) Effect of broodstock nutrition on reproductive performance of fish. *Aquaculture* 197: 25-42.
  16. Chong ASC, Saraitul DI, Zulfaizuddin O, Roshada H (2004) Effect of dietary protein level on the reproductive performance of female swordtails *Xiphophorus helleri* (Poeciliidae). *Aquaculture* 234: 381-392.
  17. Khan MA, Jafri AK, Chadha NK (2005) Effects of varying dietary protein levels on growth, reproductive performance, body and egg composition of rohu, *Labeo rohita* (Hamilton). *Aquaculture Nutrition* 11: 11-17.
  18. Gonzalez HR, Uilo MG, Liamas AH, Villarreal H (2006) Effect of dietary protein level on spawning and egg quality of redclaw crayfish *Cherax quadricarinatus*. *Aquaculture* 257: 412-419.
  19. Sink TD, Rebecca TL, Camilo P, Alejandro B, Delbert G (2010) Effects of dietary protein source and protein-lipid source interaction on channel catfish (*Ictalurus punctatus*) egg biochemical composition, egg production and quality, and fry hatching percentage and performance. *Aquaculture* 298: 251-259.
  20. Coldebella IJ, Radünz JN, Mallmann CA, Veiverberg CA, Bergamin GT, et al. (2011) The effects of different protein levels in the diet on reproductive indexes of *Rhamdia quelen* females. *Aquaculture* 312: 137-144.
  21. Johnston AT, Wiegand MD, Pronyk RJ, Dyal SD, Watchorn KE, et al. (2007) Hatching success of walleye embryos in relation to maternal and ova characteristics. *Ecology of Freshwater Fish* 16: 295-306.
  22. Syandri H (1997) The development of oocytes and testes of Bilih (*Mystacoleucus padangensis* Bleeker) in Singkarak Lake. *Journal of Fisheries Garing* 2: 1-8.
  23. De Silva SS, Radampola K (1990) Effect of dietary protein level on the reproductive performance of *Oreochromis niloticus* (L.). In: Hirano R, Hanyu I (eds.) *The Second Asian Fisheries Forum*. Asian Fisheries Society, Manila, pp. 559-564.
  24. Shim KF, Landesman L, Lam TJ (1989) Effect of dietary protein on growth, ovarian development and fecundity in the dwarf gourami, *Co&a lalia* (Hamilton). *J. Aquacult. Trop* 4: 111-123.
  25. Bhujel RC, David CL, Amjad H (2007) Reproductive performance and the growth of pre-stunted and normal Nile tilapia (*Oreochromis niloticus*) broodfish at varying feeding rates. *Aquaculture* 273: 71-79.
  26. Brion F, Tyler CR, Palazzi X, Laillet B, Porcher JM, et al. (2004) Impacts of 17 $\beta$ -estradiol, including environmentally relevant concentrations, on reproduction after exposure during embryo-larval-, juvenile- and adult-life stages in zebrafish (*Danio rerio*). *Aquatic Toxicology* 68: 193-217.
  27. Imai S, Koyama J, Fujii K (2005) Effects of 17 $\beta$ -estradiol on the reproduction of Java-medaka (*Oryzias javanicus*), a new test fish species. *Marine Pollution Bulletin* 51: 708-714.
  28. Wang HP, Gao Z, Beres B, Ottobre J, Wallat G, et al. (2008) Effects of estradiol-17 $\beta$  on survival, growth performance, sex reversal and gonadal structure of bluegill sunfish *Lepomis macrochirus*. *Aquaculture* 285: 216-223.
  29. Trijoko (2006) The effect of implantation LHRH-a hormone on gonade development of Humpback graupper (*Cromileptes altivilis*). *Journal of Fisheries Sciences* 8: 68-73.
  30. Syandri H, Azrita, Aryani N (2009) Testing the use of LHRHa to ganads maturation of bujuk (*Channa pleurothalmus*). *Jurnal Sigmatek* 3:16-23.
  31. Schreck CB, Sanchez WC, Filpatrick MS (2001) Effect of stress on reproduction gamete quality and progeny. *Aquaculture* 97: 3-24.

**Citation:** Aryani N, Suharman I (2015) Effect of Dietary Protein Level on the Reproductive Performance of Female of Green Catfish (*Hemibagrus nemurus* Bagridae). J Aquac Res Development 6: 377. doi:10.4172/2155-9546.1000377

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>

# Effect of dietary protein level on the reproductive performance of female of Green catfish ( *Hemibagrus nemurus* Bagridae )

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

15%

INTERNET SOURCES

16%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

|   |   |    |
|---|---|----|
| 1 | <a href="http://media.neliti.com">media.neliti.com</a><br>Internet Source   | 2% |
| 2 | Submitted to Jabatan Pendidikan Politeknik Dan Kolej Komuniti<br>Student Paper  | 2% |
| 3 | Submitted to Universiti Sains Malaysia<br>Student Paper   | 2% |
| 4 | Shoko Imai, Jiro Koyama, Kazunori Fujii.<br>"Effects of 17 $\beta$ -estradiol on the reproduction of Java-medaka ( <i>Oryzias javanicus</i> ), a new test fish species", Marine Pollution Bulletin, 2005<br>Publication | 2% |
| 5 | Submitted to National Penghu University of Science and Technology<br>Student Paper  | 1% |
| 6 | <a href="http://wmbc.olsztyn.pl">wmbc.olsztyn.pl</a><br>Internet Source   | 1% |
| 7 | Shubhadeep Ghosh. "Effect of probiotic on reproductive performance in female  | 1% |

livebearing ornamental fish", Aquaculture  
Research, 3/2007

Publication

---

|    |  |     |
|----|--|-----|
| 8  | <a href="http://fapet.ipb.ac.id">fapet.ipb.ac.id</a><br>Internet Source  | 1 % |
| 9  | <a href="http://goo.gl">goo.gl</a><br>Internet Source  | 1 % |
| 10 | <a href="http://www.ejpau.media.pl">www.ejpau.media.pl</a><br>Internet Source  | 1 % |
| 11 | Alexander Chong. "Assessment of soybean meal in diets for discus ( <i>Symphysodon aequifasciata</i> HECKEL) farming through a fishmeal replacement study", Aquaculture Research, 9/2003<br>Publication   | 1 % |
| 12 | Usha Jyoti Maji, Sriprakash Mohanty, Avinash Pradhan, Nikhil Kumar Maiti. "Immune modulation, disease resistance and growth performance of Indian farmed carp, <i>Labeo rohita</i> (Hamilton), in response to dietary consortium of putative lactic acid bacteria", Aquaculture International, 2017<br>Publication | 1 % |
| 13 | El-Sayed, A.F.M.. "Effects of dietary lipid source on spawning performance of Nile tilapia ( <i>Oreochromis niloticus</i> ) broodstock   | 1 % |



reared at different water salinities",  
Aquaculture, 20050729

Publication

14

El-Sayed, A.F.M.. "Effects of dietary protein and energy levels on spawning performance of Nile tilapia (*Oreochromis niloticus*) broodstock in a recycling system",  
Aquaculture, 20080801

Publication

1 %

15

[netlib.org](http://netlib.org)

Internet Source

1 %

16

[pinpdf.com](http://pinpdf.com)

Internet Source

<1 %

17

[almasdi.staff.unri.ac.id](http://almasdi.staff.unri.ac.id)

Internet Source

<1 %

18

Fábio P. Arantes, Hélio B. Santos, Elizete Rizzo, Yoshimi Sato, Nilo Bazzoli. "Influence of water temperature on induced reproduction by hypophysation, sex steroids concentrations and final oocyte maturation of the "curimatã-pacu" *Prochilodus argenteus* (Pisces: Prochilodontidae)", *General and Comparative Endocrinology*, 2011

Publication

<1 %

19

Muchlisin Zainal Abidin. "Influence of dietary protein levels on growth and egg quality in broodstock female bagrid catfish (*Mystus*

<1 %

nemurus Cuv. & Val.)", Aquaculture Research, 3/2006

Publication

20

[evols.library.manoa.hawaii.edu](http://evols.library.manoa.hawaii.edu)

Internet Source

<1 %

21

Tu, Yongqin, Shouqi Xie, Dong Han, Yunxia Yang, Junyan Jin, Haokun Liu, and Xiaoming Zhu. "Growth performance, digestive enzyme, transaminase and GH-IGF-I axis gene responsiveness to different dietary protein levels in broodstock allogonogynetic gibel carp (*Carassius auratus gibelio*) CAS III", Aquaculture, 2015.

Publication

<1 %

22

EBRAHIMI, Ghaffar and OURAJI, Hossein. "Growth performance and body composition of kutum fingerlings, *Rutilus frisii kutum* (Kamenskii 1901), in response to dietary protein levels", TÜBİTAK, 2012.

Publication

<1 %

23

[researchonline.jcu.edu.au](http://researchonline.jcu.edu.au)

Internet Source

<1 %

24

AGUS KURNIA, SHUICHI SATOH, SATOSHI HANZAWA. "Effect of *Paracoccus* sp. and their Genetically Modified on Skin Coloration of Red Sea Bream", HAYATI Journal of Biosciences, 2010

Publication

<1 %

|    |   |      |
|----|---|------|
| 25 | <a href="http://link.springer.com">link.springer.com</a><br>Internet Source   | <1 % |
| 26 | <a href="http://webdoc.sub.gwdg.de">webdoc.sub.gwdg.de</a><br>Internet Source   | <1 % |
| 27 | <a href="http://www.preprints.org">www.preprints.org</a><br>Internet Source   | <1 % |
| 28 | <a href="http://zenodo.org">zenodo.org</a><br>Internet Source   | <1 % |
| 29 | "Sex Control in Aquaculture", Wiley, 2018<br>Publication  | <1 % |
| 30 | Ghanawi, Joly, and I. Patrick Saoud. "Molting, reproductive biology, and hatchery management of redclaw crayfish <i>Cherax quadricarinatus</i> (von Martens 1868)", Aquaculture, 2012.<br>Publication   | <1 % |
| 31 | J.Y. LI, Z.L. GUO, X.H. GAN, D.L. WANG, M.F. ZHANG, Y.L. ZHAO. "Effect of different dietary lipid sources on growth and gonad maturation of pre-adult female <i>Cherax quadricarinatus</i> (von Martens)", Aquaculture Nutrition, 2011<br>Publication | <1 % |
| 32 | Jaya-Ram, A.. "Influence of dietary HUFA levels on reproductive performance, tissue fatty acid profile and desaturase and elongase mRNAs  | <1 % |

expression in female zebrafish *Danio rerio*",  
Aquaculture, 20080603

Publication

---

33

Nooshin Mehdinejad, Mohammad Reza Imanpour, Valiollah Jafari. "Combined or Individual Effects of Dietary Probiotic, *Pediococcus acidilactici* and Nucleotide on Reproductive Performance in Goldfish (*Carassius auratus*)", Probiotics and Antimicrobial Proteins, 2018

Publication

---

34

Santinha. "Effects of the dietary protein : lipid ratio on growth and nutrient utilization in gilthead seabream (*Sparus aurata* L.)", Aquaculture Nutrition, 8/1999

Publication

---

35

[revistamvz.unicordoba.edu.co](http://revistamvz.unicordoba.edu.co)

Internet Source

---

36

[ricerivers.vcu.edu](http://ricerivers.vcu.edu)

Internet Source

---

37

Hipólito Fernández-Palacios, Birgitta Norberg, Marisol Izquierdo, Kristin Hamre. "Effects of Broodstock Diet on Eggs and Larvae", Wiley, 2011

Publication

---

38

Watanabe, T.. "Broodstock nutrition research on marine finfish in Japan", Aquaculture,

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

20031110

Publication

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On