

Evaluation of salted trash fish as a protein source replacing fishmeal in the diet for river catfish (*Hemibagrus nemurus*)

by Indra Suharman


Submission date: 07-Apr-2023 01:25AM (UTC+0700)

Submission ID: 2057771473

File name: AACL_bioflux_vol_9-3_2016.pdf (89.69K)

Word count: 5726

Character count: 28007



8

Evaluation of salted trash fish as a protein source replacing fishmeal in the diet for river catfish (*Hemibagrus nemurus*)

Bustari Hasan, Iskandar Putra, Indra Suharman, Dian Iriani

3

Faculty of Fisheries and Marine Sciences, Universitas Riau, Jl. HR. Subratas Km 12.5, Pekanbaru, Indonesia. Corresponding author: B. Hasan, bustarih@yahoo.com

Abstract. This research was conducted to evaluate the effect of substitution of fishmeal by salted trash fish in the diet on growth, feed efficiency, nutrient utilization and body composition of river catfish (*Hemibagrus nemurus*). Five formulated diets were prepared to contain 38% protein and 3.25 kcal g⁻¹ digestible energy (DE) as an optimum dietary protein and energy for the fish. One diet was control diet which contained fishmeal without salted trash fish (FM), and other diets were trash fish diets which fishmeal was reduced and proportionally replaced by salted trash fish 25% (STF-25), 50% (STF-50), 75% (STF-75) and 100% (STF-100). Catfish fingerlings, ±5 g size obtained from local fish hatchery were stocked into triplicate 2 x 2 x 1.5 m floating net cages at a density of 100 fish cage⁻¹, and fed experimental diets at satiation, twice a day for 8 weeks. Results indicated that the salted trash fish diets were characterized by higher salt, ash and fiber but similar in amino acid profiles as well as water stability as compared to control diet. The survival rate was reached 100% in all treatments. Inclusion of salted trash fish at level 25% in the diet increased weight gain as compared to control diet; and inclusion up to 50% did not affect weight gain, specific growth rate, food consumption, food efficiency, protein efficiency and protein retention; but further increase reduced weight gain, specific growth rate, food efficiency, protein efficiency and protein retention. Salted trash fish diets did not affect fish body protein and ash, but increased body moisture and decreased body fat. Essential amino acid profiles of the fish fed salted trash fish diets were similar to that fish fed control diet. It was concluded that the salted trash fish could be included in the river catfish diet up to 50% as replacement for 50% fishmeal without negative effect on weight gain, specific growth rate, food consumption, food efficiency, protein efficiency, protein retention and proximate body composition of river catfish.

Key Words: fishmeal diet, trash fish diet, river catfish, growth performance, fish body composition.

Introduction. River catfish (*Hemibagrus nemurus* Valenciennes, 1840) is a popular and demanded river catfish species for both fresh and smoked fish consumption in Riau Province, Indonesia. Naturally, the fish was caught from the rivers, lakes and reservoir, however, its population in the wild was decreasing due to overfishing and environmental damage, therefore, the next supply of the fish would depend on aquaculture production. The culture of the river catfish now has been developed; artificial breeding and aquaculture techniques have been available; however, the fish was required relative high dietary protein (approximately 38-42%) for its optimum growth and high flesh quality (Khan et al 1993; Hasan et al 1999; Hasan et al 2013).

Fishmeal (FM) is the most preferable and digestible protein source in the diet for most farmed fish including river catfish due to its amino acid, fatty acid, energy, vitamins and minerals balance (Tacon 1993; Eguia 1998; Khan et al 1993; Hasan et al 1999; 2013; Abdelghany 2003). However, its price is high and its production is limited and scarcely available, import dependent, hence it is not economical for small scale feed manufacturer. Since the demand for FM continues to grow, while its production is expected to remain constant (New & Wijkstrom 2007; Borgeson et al 2006), more attention must be paid to utilizing local material as an alternative protein source replacing FM in the fish diet.

Trash fish from marine by-catch is a potential protein source as a replacement for conventional FM in the fish diet due to its high protein content, averaging 66.1% dry

weight (Li et al 2004) and its abundant supply which is estimated more than 4.39 million tons or 62.6% of the total annual marine catches in Indonesia (Davies et al 2009; Anon 2010). However, most trash fish production is scattered at small scale fishermen in remote fishing ground. The fish is highly perishable while refrigeration facilities and ice are expensive and scarcely available, thus an economic and practical preservation method is required to collect and transport it to feed processing industries.

Salting is a cheap and practical method for trash fish preservation as the salt is easily available and the salting process only needs simple technology. In Riau Province, Indonesia, salted trash fish has been used for protein source in the diets for some farm raised fish; and its acceptability by fish was reported to vary with fish species and salt concentration (personal communication). Our personal observation showed that farmers included salted trash fish as protein source up to 30% in the diet for some farm raised fish such as pangasius catfish (*Pangasius hypophthalmus*) and tilapia (*Oreochromis niloticus*); but there was no scientific information on its effect on growth and fish body composition, especially for river catfish. Hence, this search was to evaluate the effect of substitution of dietary FM by salted trash on growth performance and fish body composition of river catfish.

Material and Method

Diet preparation and analysis. Dried salted trash fish of marine by catch was taken from a local feed stuff supplier. The trash fish was desalted in boiling water (1 part of trash fish: 2 parts water) for 15 minutes and pressed to reduce its salt concentration, dried and ground. Five diets were formulated to contain 38% protein and 3.25 kcal g⁻¹ digestible energy (DE), which the levels were within the range of dietary protein and energy requirement optimum growth of river catfish, 34-42% protein and 2.75-3.25 kcal g⁻¹ energy (Khan et al 1993; Hasan et al 1999; Hasan et al 2013; Hasan et al 2016). One diet was control diet containing conventional FM without salted trash fish; and other diets were salted trash fish diets which FM was reduced and proportionally replaced with salted trash fish 25% (STF-25), 50% (STF-50), 75% (STF-75) and 100% (STF-100), respectively.

The diets were pelleted with extruder machine, dried and analyzed for salt concentration, proximate and amino acid composition as well as stability in the water. Salt concentration and proximate composition analyses were conducted based on AOAC method (AOAC 1990). Salt concentration was analyzed by titration with AgNO₃ 0.1 N after incineration at 500°C. Moisture was determined after the sample was oven-dried at 105°C until constant in weight. Ash was determined after the sample was incinerated at 500°C for 5 hours. Protein was analyzed by micro-Kjeldahl procedure and crude protein was estimated as N x 6.25. Crude fat was determined after the sample was Soxhlet-extracted with petroleum ether. Amino acids analyses were conducted by HPLC using Pico-tag method (Waters USA) according to Cohen et al (1989). Amino acid profile was determined after hydrolyzed under nitrogen in 6N HCl at 110°C for 24 hours and the amino acids were calculated in percent protein.

Pellet stability in the water was determined based on Fagbenro & Jauncey (1995). Triplicate 50 g samples of pellet diets were placed on sieve and slowly immersed in 40L aquarium containing distilled water at room temperature for 10 and 30 minutes. The sieves were removed to dry for 1 minute, oven-dried at 105°C for 2 hours, cooled in a desiccator and reweighed. Water stability was calculated as the percentage difference in weight after reweighing and expressed as percentage loss of dry matter (% LDM).

Feeding trial. River catfish fingerlings, ±5 g in size were taken from a local fish hatchery in Kampar district. The fish was stocked in triplicate 2 x 2 x 1.5 m experimental floating net cages at stocking density of 100 fish per cage; and experimental fish fed the formulated diets for 8 weeks, from June to August 2015. The fish were fed at satiation twice daily, 9.00 and 16.00. All the fish were weighed at the beginning and two week intervals. Water quality (temperature and dissolved oxygen) and fish mortality were recorded daily. Twenty fish at the beginning of experiment and all fish of each cages at

the end of the experiment were taken randomly and analyzed for proximate and amino acid compositions. Growth parameters were calculated by following formula:

$$\begin{aligned} \text{Weight gain (g fish}^{-1}\text{)} &= (\text{Final fish weight} - \text{Initial fish weight}) \\ \text{Specific growth rate (\% day}^{-1}\text{)} &= [(\text{Ln final weight} - \text{Ln initial weight}) / \text{days of trial}] \times 100; \\ \text{Survival rate (\%)} &= (\text{Total final survived fish}) / (\text{Total initial fish}) \times 100; \\ \text{Food consumption (g fish}^{-1}\text{)} &= (\text{Total food consumed fish}^{-1}); \\ \text{Food efficiency ratio} &= (\text{Weight gain, g}) / (\text{Food consumed, g}); \\ \text{Protein efficiency ratio} &= (\text{Weight gain, g}) / (\text{Protein consumed, g}); \\ \text{Protein retention} &= (\text{Protein gain, g}) / (\text{Protein consumed, g}) \times 100 \end{aligned}$$

Data analysis. Triplicate data were analyzed by the Analysis of Variance using SPSS (SPSS 2000). Least significant different test was used to determine the differences among individual means.

Results and Discussion

Feed ingredients, formulation and proximate composition of the diets. Proximate and salt composition of salted trash fish and other feed ingredients used in the experimental diets was presented in Table 1. Salted trash fish contained 55.29% protein, 6.35% fat, 9.76% moisture, 14.67% ash, 2.30% crude fiber, 11.93% nitrogen free extract (NFE) and 10.19% salt. Protein composition of salted trash fish was similar to conventional FM, but its fat content was lower; and ash, NFE as well as salt content was higher. Lower fat in salted trash fish in this study may be due to boiling and pressing effect as the salted trash fish before inclusion to the diets was boiled in boiling water and pressed for reducing its salt concentration (desalting). During boiling and pressing process, the trash fish fat may be washed, so its fat content reduced.

Proximate composition and salt concentration of the feed ingredients

Table 1

Feed ingredients	Proximate composition (%)						
	Protein	Fat	Moisture	Ash	Fiber	NFE*	Salt
Fishmeal	56.68	13.97	8.48	12.24	2.00	6.63	2.84
Salted trash fish	55.29	6.35	9.76	14.67	2.30	11.93	10.19
Soybean meal	40.50	17.69	9.90	8.33	8.50	15.08	-
Rice bran	8.54	6.21	6.83	12.52	15.00	50.90	-

*NFE calculated as 100- (protein, fat, moisture, ash, fiber).

Formulation and proximate composition of the experimental diets was shown in Table 2. Proportions of each diet components were made similar, except for conventional FM and salted trash fish. Protein and energy composition of each diet was maintained 38% and 3.25 kcal g⁻¹ DE respectively as dietary requirement for optimum growth of fish; and palm oil was used to maintain energy balance in each diet. Proximate compositions of salted trash fish diets were similar to control diet, except salt concentration which was higher in salted trash fish diets than that for control diet. Salt concentration of FM, STF-25, STF-50, STF-75 and STF-100 diet was 0.96%; 1.63%; 2.27%; 2.97% and 3.72% respectively; and their concentrations increased as the increasing amount of salted trash fish in the diets. Essential amino acid profiles of salted trash fish diets were similar to that for control diet.

Essential amino acid profiles of salted trash fish and control diet were similar (Table 3). The A/E ratio (% essential amino acid content/total amino acid) of river catfish which is considered as an indicator for amino acid balance in the fish diets (Arai 1981; Ogata et al 1983; Murai et al 1984; Khan et al 2013; Hasan et al 2001) was also similar between trash fish diets and control diet. As proximate composition and amino acid profile of the salted trash fish diets and control diet was similar, salt concentration

therefore was the main factor characterizing salted trash fish diets different from control diet.

50

Table 2

Proximate and salt concentration of the experimental diets

Ingredients	Diets (%)				
	FM	STF-25	STF-50	STF-75	STF-100
Fishmeal	35	26.25	17.5	8.75	0
Salted trash fish	0	8.75	17.5	26.25	35
Soybean meal	30	30	30	30	30.3
Rice bran	30.5	30.5	29.5	29.2	28
Palm oil	3.5	3.5	4.5	4.8	5.7
Vitamin and mineral mix ^a	1	1	1	1	1
Proximate composition by analysis					
Dry matter	91.26	91.95	91.62	91.58	91.95
Protein	38.12	37.59	37.43	37.72	37.21
Fat	16.76	16.65	16.49	16.52	16.51
Fiber	8.92	9.77	9.86	9.92	10.38
Ash	12.76	13.17	13.27	13.52	13.62
NFE ^b	23.44	22.82	22.95	22.32	22.28
Energy (Kcal ^{g-1} DE)	3.27	3.23	3.21	3.21	3.20
Salt	0.96	1.63	2.27	2.97	3.72

^aVitamin and mineral mix: Vit A, 2750 IU; Vit D, 550,000 IU; Vit E, 25,000 IU; Vit K, 5,000 mg; Choline, 250,000 mg; Niacin, 50,000 mg; Riboflavin, 10,000 mg; Pyridoxine, 10,000 mg; Calcium D-pantothenate, 25,000 mg; Biotin, 50 mg; Folic acid, 2,500 mg; Cyanocobalamin, 10 mg; Ascorbic acid, 50,000 mg; K₂HPO₄, 30%; KCl, 8.4%; MgSO₄, 14.8%; CaHPO₄·2H₂O, 27.4%; FeCl₃, 1.4%; MnSO₄·7H₂O, 0.2%; CaCO₃, 16.8%; ^bNFE calculated as 100- (protein, fat, moisture, ash, fiber).

20

Table 3

A/E ratio (% essential amino acid/total essential amino acid) of experimental diets and *Hemibagrus nemurus* (% protein)

Essential amino acid	Diets					<i>Hemibagrus nemurus</i>
	FM	STF-25	STF-50	STF-75	STF-100	
Arginine	14.47	14.37	14.69	14.62	14.79	14.37
Histidine	5.45	5.35	5.42	5.37	5.42	5.56
Isoleucine	7.22	6.98	7.12	6.78	6.74	6.94
Leucine	15.20	14.95	15.14	14.90	14.95	13.80
Lysine	14.49	14.19	14.10	14.07	14.03	14.08
Methionine ^a	7.17	6.98	6.97	7.06	7.25	7.13
Phenylalanine ^b	15.24	15.99	15.34	16.24	16.10	15.66
Threonine	8.91	8.69	8.72	8.66	8.21	8.83
Valine	11.85	12.60	12.70	12.46	12.79	12.05
Tryptophan	ND ^c	ND	ND	ND	ND	ND

^aMethionine + Cystine, ^bPhenylalanine + Tyrosine, ^cND: Not determined.

30

Water stability of pelleted diets. Water stability of experimental pelleted feed diets was presented in Table 4. The water stability of pelleted feed diets was measured by percentage loss of dry matter (LDM) after 10 and 30 minutes in the water, and the best water stability was the feed with a minimum loss in dry matter. The LDM values in this study were not different between salted trash fish pelleted diets and control pelleted diet ($p > 0.05$), indicating that there was no inclusion effect of salted trash fish in the diet on water stability of the pelleted feed. LDM values of control diet and salted trash fish diets were 2.02-2.04% for 10 minutes and 6.37-6.38% for 30 minutes. These values were considered very stable as a minimum LDM value for catfish feed is < 10% for 5 minutes (Wood et al 1985; Fagbenro & Jauncey 1995).

Table 4

Water stability of feed (% Loss of Dry Matter, LDM) for 10 and 30 minutes

Diets (pellet)	Water stability (% LDM)	
	LDM-10 Minutes	LDM-30 Minutes
FM	2.04±0.07 ^a	6.38±0.07 ^a
STF-25	2.02±0.13 ^a	6.38±0.05 ^a
STF-50	2.03±0.05 ^a	6.37±0.05 ^a
STF-75	2.03±0.08 ^a	6.37±0.06 ^a
STF-100	2.04±0.07 ^a	6.38±0.04 ^a

Note: Means in the same column with the same superscript were not different ($p < 0.05$).

Water quality management. Water quality values along the feeding trial were shown in Table 5. Dissolved oxygen in the morning and afternoon was 3.40-5.50 mg L⁻¹ and 3.40-5.50 mg L⁻¹ respectively; temperature was 28-28.9°C and 31-33.3°C respectively; and pH was 6.9-7.3 and 6.9-7.3, respectively. The values were within the acceptable ranges for the culture of warm water fishes. Dissolved oxygen which is an important parameter affecting growth responses of fish is acceptable since the recommended value for growth of catfish is greater than 3 mg L⁻¹ (Weeks & Ogburn 1973).

Table 5

Water quality during feeding trial

Parameter	Water quality	
	Morning	Afternoon
Dissolved oxygen (mg L ⁻¹)	3.40-5.50	3.40-5.50
Temperature (°C)	28.0-28.9	31.0-33.3
pH	6.9-7.3	6.9-7.3

Growth performance and food utilization. The inclusion effect of salted trash fish as a replacement for conventional FM in the diet on survival rate, weight gain and specific growth rate were shown in Table 6; and food consumption, food efficiency ratio, protein efficiency ratio and protein retention in Table 7. Survival rate of the fish was 100%; and there was no difference in survival rate among the fish fed experimental diets ($p > 0.05$). Inclusion of salted trash fish at a level of 25% in the diet (STF-25) increased weight gain compared to control diet ($p < 0.05$); and inclusion up to 50% (STF-50) did not affect weight gain and specific growth rate ($p > 0.05$); however, inclusion over 50% decreased weight gain and specific growth rate ($p < 0.05$). Survival rate of the fish was 100%; and there was no difference in survival rate among the fish fed experimental diets ($p > 0.05$). Inclusion of salted trash fish at a level of 25% in the diet (STF-25) increased weight gain compared to control diet ($p < 0.05$); and inclusion up to 50% (STF-50) did not affect weight gain, specific growth rate, food conversion, protein retention and survival rate ($p < 0.05$). These indicated that the salted trash fish diets were acceptable and utilizable by the fish, and could be included in the fish diet replacing FM up to 50% without adverse effect on growth, food conversion and utilization. The fact that at a low level of salted trash fish in the diet increased food consumption and weight gain was probably due to salt concentration of the diet which at a small amount (1.63%) enhanced palatability of the diets.

However, the fact that the inclusion of salted trash fish > 50% replacing FM in the diet reduced weight gain, specific growth rate, food conversion and protein retention ($p < 0.05$) might be due to higher concentration of salt in the diets (> 2.27%) which reduced palatability of the diet, thus decreased food consumption, weight gain, food efficiency and nutrient utilization. Reduced growth due to the excess amount of salt in the fish diets was also reported in juvenile of flounder (*Paralichthys olivaceus*) by Park et al (2000).

63 Table 6
The effect of salted trash fish diets on survival rate, weight gain, specific growth rate

Diet	Initial weight (g)	Final weight (g)	Survival rate (%)	Weight gain (%)	Specific growth rate (%)
FM	5.16	21.52	100	16.36 ^c	2.39 ^c
STF-25	5.25	25.65	100	20.41 ^d	2.65 ^{b^c}
STF-50	5.29	20.17	100	14.88 ^{bc}	2.22 ^{abc}
STF-75	4.98	16.48	100	11.50 ^{ab}	1.99 ^{ab}
STF-100	4.78	15.47	100	10.49 ^a	1.92 ^a

Note: Means in the same column with the same superscript were not different ($p < 0.05$).

Table 7
The effect of salted trash fish diets on food consumption, food efficiency ratio, protein efficiency ratio and protein retention

Diets	Food consumption (g/fish)	Food efficiency ratio	Protein efficiency ratio	Protein retention
FM	35.88 ^{bc}	0.45 ^a	1.20 ^b	15.94 ^b
STF-25	45.33 ^d	0.45 ^a	1.18 ^b	15.90 ^b
STF-50	36.03 ^{bc}	0.41 ^a	1.08 ^{ab}	13.82 ^b
STF-75	32.56 ^a	0.35 ^b	0.93 ^a	11.71 ^a
STF-100	31.98 ^a	0.33 ^b	0.86 ^a	10.42 ^a

Note: Means in the same column with the same superscript were not different ($p < 0.05$).

1
The effect of salted trash fish diets on proximate and amino acid composition of fish body was presented in Table 8. There was no effect of salted trash fish diets on protein and ash body composition of fish as compared to control diet ($p > 0.05$); however, moisture increased and fat decreased as the salted trash fish increase in the diet ($p < 0.05$). Amino acid profile of whole body fish was similar between fish fed control and salted fish diets. The inclusion effects of alternative animal and plant protein sources for FM in the diets on proximate body composition of fish has been reported in channel catfish (*Ictalurus punctatus*) (Peterson et al 2012), Nile tilapia (*Oreochromis niloticus*) (Labib et al 2012); olive flounder (*Paralichthys olivaceus*) (Deng et al 2006; Uy et al 2006; Kikuchi 1999; Lee et al 2012; Kader & Koshio 2012; Kim et al 2014); Atlantic salmon (*Salmo salar*) (Pratoomyot et al 2010; Pratoomyot et al 2011), rainbow trout (*Oncorhynchus mykiss*) (Lee et al 2010); black sea bream (*Sparus macrocephalus*) (Zhou et al 2011); coho salmon (*Oncorhynchus kisutch*) (Watson et al 2014), large yellow croaker (*Larimichthys croceus*) (Yi et al 2015); California yellowtail (*Seriola lalandi*) (Suentello et al 2015). In the present study, inclusion of salted trash fish in the diets did not affect protein and ash body composition of fish; however, fish body fat tended to decrease and body moisture increase as the increasing of the salted trash fish levels in the diets. Moreover, reduced food consumption as the level of dietary FM excessively decreased had also been identified as a factor reducing growth and specific growth rate in Atlantic salmon (Pratoomyot et al 2010; Pratoomyot et al 2011).

The effect of salted trash fish diets on proximate composition and amino acid profiles of harvested fish

Proximate composition	Diets (%)				
	FM	STF-25	STF-50	STF-75	STF-100
Moisture	72.42 ^a	73.29 ^b	74.44 ^c	74.76 ^c	74.80 ^c
Ash	2.91 ^a	3.04 ^a	3.10 ^a	3.25 ^a	3.29 ^a
Protein	13.39 ^a	13.84 ^a	13.43 ^a	13.26 ^a	13.22 ^a
Fat	8.21 ^d	7.59 ^c	6.22 ^b	5.69 ^a	5.51 ^a
Amino acid (% protein)	FM	STF-25	STF-50	STF-75	STF-100
Aspartic acid	9.44	9.42	8.19	8.08	7.38
Cystine	0.14	0.12	0.19	0.18	0.13
Glutamic acid	13.77	14.04	13.06	13.20	13.13
Serine	2.91	3.07	2.79	2.84	3.02
Glycine	5.31	5.69	5.63	5.58	5.64
Histidine	1.62	1.51	1.38	1.24	1.09
Arginine	4.27	4.25	3.96	3.98	3.89
Threonine	2.63	2.67	2.48	2.35	2.13
Alanine	5.54	5.60	5.12	5.15	5.26
Proline	3.37	3.36	3.12	3.25	3.90
Tyrosine	2.10	2.26	2.09	2.11	2.39
Valine	3.68	3.76	3.46	3.57	3.38
Methionine	2.34	2.42	2.18	2.20	2.05
Isoleucine	3.27	3.36	2.95	2.94	2.74
Leucine	4.50	4.86	4.14	4.24	3.63
Phenylalanine	5.38	5.37	5.00	5.04	4.78
Lysine	4.14	4.21	3.84	3.81	3.65
Total	74.42	75.98	69.59	69.76	68.21

Note: Means in the same row with the same superscript were not different ($p < 0.05$).

Conclusions. Trash fish diets were characterized by higher salt but similar in amino acid profile and water stability as compared to control diets. Inclusion of salted trash fish 25% in the diet increased weight gain and specific growth rate, and inclusion up to 50% did not affect weight gain, specific growth rate, food conversion, protein retention and survival rate as compared to control diet; but inclusion more than 50% reduced food conversion and protein retention. Salted trash fish diets did not affect fish body protein and ash, but increased body moisture and decreased body fat. There was no marked difference in essential amino acid profiles between the fish fed salted trash fish diets and control diet. The salted trash fish therefore could be included in the river catfish diet up to 50% as replacement for 50% FM without negative effect on growth and body composition.

Acknowledgements. The authors wish to express their grateful acknowledgement to University of Riau for providing research grant. Appreciations are also extended to Rahmad dan Amsar, the students at Faculty of Fisheries and Marine Science, University of Riau Pekanbaru for field technical assistance and Mr. Syafrudin, a directeur of fish feed processing plant, Kampar for feed pelleting processing.

References

- Abdelghany A. E., 2003 Partial and complete replacement of fishmeal with gambusia meal in diets for red tilapia (*Oreochromis niloticus* x *O. mossambicus*). *Aquaculture Nutrition* 9:145-154.
- Anon, 2010 National report on by-catch management and reduction of discard. National Report Indonesia, Rebyc II CTI, Jakarta, May 2010.

- AOAC, 1990 Official Method of Analysis. The Association of Official Analytical Chemist, Washington DC, USA, 1094 pp.
- Arai S., 1981 A purified test diet for coho salmon (*Oncorhynchus kisutch*) fry. Bulletin of the Japan Society of Scientific Fisheries 47:547-550.
- Buentello A., Jirsa D., Barrows F. T., Drawbridge M., 2015 Minimizing fishmeal use in juvenile California yellowtail (*Seriola lalandi*) diets using non-GM soybeans selectively bred for aquafeeds. Aquaculture 435:403-411.
- Borgeson T. L., Racz V. J., Wilkie D. C., White L. J., Drew M. D., 2006 Effect of replacing fishmeal and oil with simple or complex mixtures of vegetable ingredients in diets fed to Nile tilapia (*Oreochromis niloticus*). Aquaculture Nutrition 12:141-149.
- Cohen S. A., Meys M., Tarvin T. L., 1989 The PicoTag method. A manual of advanced techniques for amino acid analysis. Waters Chromatography Division, 34 Maple Street, Milford, MA, pp. 1-12.
- Davies R. W. D., Cripps S. J., Nickson A., Porter G., 2009 Defining and estimating global marine fisheries bycatch. Marine Policy (2009), doi:10.1016/j.marpol.2009.01.003. 29 pp.
- Deng J., Mai K., Ai Q., Zhang W., Wang X., Xu W., Liufu Z., 2006 Effects of replacing fish meal with soy protein concentrate on feed intake and growth of juvenile Japanese flounder (*Paralichthys olivaceus*). Aquaculture 258:503-513.
- Eguia P. L. V., 1998 Development of artificial diet and optimum feeding strategy for Malaysian catfish (*Mystus nemurus*) larvae. Master thesis, University Putra Malaysia, Serdang, Malaysia, 107 pp.
- Fagbenro O. A., Jauncey K., 1995 Growth and protein utilization by juvenile catfish (*Clarias gariepinus*) fed dry diets containing co-dried lactic acid fermented fish silage and protein feedstuffs. Bioresource Technology 51:29-35.
- Hasan B., Saad C. R., Alimon A. A., Kamarudin M. S., Hassan Z., 1999 The effects of various dietary energy levels and protein concentrations on growth performance of river catfish (*Mystus nemurus*). Malaysian Journal of Animal Science 5(1-2):83-92.
- Hasan B., Saad C. R., Alimon A. R., Kamarudin M. S., Hassan Z., 2001 Replacement of fishmeal with co-dried fish silage in the diet for *Mystus nemurus*. Malaysian Journal of Animal Science 7(1):69-79.
- Hasan B., Suharman I., Desmelati, Iriani D., 2013 [An improvement of carcass quality characteristic of farmed baung catfish (*Mystus nemurus*) for smoked catfish processing through dietary protein and energy formulation]. Lembaga Penelitian Universitas Riau Pekanbaru, pp. 167-171. [in Indonesian]
- Hasan B., Suharman I., Desmelati D., Iriani D., 2016 Carcass quality of raw and smoked fish fillets prepared from cage raised river catfish (*Hemibagrus nemurus*, Valenciennes, 1840) fed high protein-low energy and low protein-high energy diets. Jurnal Teknologi (Science & Engineering) 78(4-2):21-25.
- Kader M. A., Koshio S., 2012 Effect of composite mixture of seafood by-products and soybean proteins in replacement of fishmeal on the performance of red sea bream (*Pagrus major*). Aquaculture 368:95-102.
- Khan M. S., Ang K. J., Ambak M. A., Saad C. R., 1993 Optimum dietary protein requirement of a Malaysian catfish (*Mystus nemurus*). Aquaculture 112:227-235.
- Kikuchi K., 1999 Use of defatted soybean meal as a substitute for fish meal in diets of Japanese flounder (*Paralichthys olivaceus*). Aquaculture 179:3-11.
- Kim H. S., Jung W. G., Myung S. H., Cho S. H., Kim D. S., 2014 Substitution effects of fishmeal with tuna byproduct meal in the diet on growth, body composition, plasma chemistry and amino acid profiles of juvenile olive flounder (*Paralichthys olivaceus*). Aquaculture 431:92-98.
- Labib E. M. H., Zaki M. A., Mabrouk, 2012 Nutritional studies on partial and total replacement of fishmeal by jojoba meal (*Simmondsia chinensis*) in Nile tilapia (*Oreochromis niloticus*) fingerlings diets. APCBEE Procedia 4:196-203.
- Lee J., Choi I. C., Kim K. T., Cho S. H., Yoo J. Y., 2012 Response of dietary substitution of fishmeal with various protein sources on growth, body composition and blood chemistry of olive flounder (*Paralichthys olivaceus*, Temminck & Schlegel, 1846). Fish Physiology and Biochemistry 38:735-744.

- Lee K. J., Powell M. S., Barrows F. T., Smiley S., Bechtel P., Hardy R. W., 2010 Evaluation of supplemental fish bone meal made from Alaska seafood processing byproducts and dicalcium phosphate in plant protein based diets for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 302:248-255.
- Li P., Wang X., Hardy R. W., Gatlin III D. M., 2004 Nutritional value of fisheries by-catch and by-product meals in the diet of red drum (*Sciaenops ocellatus*). *Aquaculture* 236:485-496.
- Murai T., Akiyama T., Nose T., 1984 Effect of amino acid balance on efficiency in utilization of diet by fingerling carp. *Bulletin of the Japan Society of Scientific Fisheries* 50:893-897.
- New M. B., Wijkstrom U. N., 2002 Use of fishmeal and fish oil in aquafeeds: further thoughts on the fishmeal trap. *FAO Fish Circ. No. 975*, FAO, Rome, Italy, 61 pp.
- Ogata H., Arai S., Nose T., 1983 Growth responses of cherry salmon (*Oncorhynchus masou*) and amago salmon (*Oncorhynchus rhodurus*) fry fed purified casein diets supplemented with amino acids. *Bulletin of the Japan Society of Scientific Fisheries* 49:1381-1385.
- Park G. S., Takeuchi T., Seikai T., Yoshinaga T., 2000 The effects of residual salts and free amino acids in mysid meal on growth of juvenile Japanese flounder (*Paralichthys olivaceus*). *Nippon Suisan Gakkaishi* 66(4):697-704.
- Peterson B. C., Booth N. J., Manning B. B., 2012 Replacement of fishmeal in juvenile channel catfish (*Ictalurus punctatus*) diets using a yeast-derived protein source: the effects on weight gain, food conversion ratio, body composition and survival of catfish challenged with *Edwardsiella ictaluri*. *Aquaculture Nutrition* 18:132-137.
- Pratoomyot J., Bendiksen E. Å., Bell J. G., Tocher D. R., 2010 Effects of increasing replacement of dietary fishmeal with plant protein sources on growth performance and body lipid composition of Atlantic salmon (*Salmo salar* L.). *Aquaculture* 305:124-132.
- Pratoomyot J., Bendiksen E. Å., Campbell P. J., Jauncey K. J., Bell J. G., Tocher D. R., 2011 Effects of different blends of protein sources as alternatives to dietary fishmeal on growth performance and body lipid composition of Atlantic salmon (*Salmo salar* L.). *Aquaculture* 316:44-52.
- SPSS 2000 SPSS for Windows, Release 10. SPSS Inc, Chicago, IL.
- Tacon A. G. J., 1993 Feed ingredients for warm water fish: fishmeal and other processed feed stuffs. *FAO Fish Circ. No. 856*, FAO, Rome, Italy, 64 pp.
- Uyan O., Koshio S., Teshima S., Ishikawa M., Thu M., Alam M. S., Michael F. R., 2006 Growth and phosphorous loading by partially replacing fishmeal with tuna muscle by-product powder in the diet of juvenile Japanese flounder (*Paralichthys olivaceus*). *Aquaculture* 257:437-445.
- Watson A. M., Buentello A., Place A. R., 2014 Partial replacement of fishmeal, poultry by-product meal and soy protein concentrate with two non-genetically modified soybean cultivars in diets for juvenile cobia (*Rachycentron canadum*). *Aquaculture* 434:129-136.
- Weeks J. P., Ogburn C. B., 1973 Catfish production guide. *Circ. E-18*. Alabama Cooperative Extension Service, Auburn University, Auburn, USA, 35 pp.
- Wood J. F., Capper B. S., Nicolaidis L., 1985 Preparation and evaluation of diet containing fish silage, cooked fish preserved with formic acid and low-temperature-dried fishmeal as protein sources for mirror carp (*Cyprinus carpio*). *Aquaculture* 44:27-40.
- Yi X., Li J., Xu W., Zhou H., Smith A. A., Zhang W., Mai K., 2015 Shrimp shell meal in diets for large yellow croaker (*Larimichthys croceus*): effects on growth, body composition, skin coloration and anti-oxidative capacity. *Aquaculture* 441:45-50.
- Zhou F., Xiao J. X., Hua Y., Ngandzali B. O., Shao Q. J., 2011 Dietary L-methionine requirement of juvenile black sea bream (*Sparus macrocephalus*) at a constant dietary cystine level. *Aquaculture Nutrition* 17:469-481.

Received: 05 January 2016. Accepted: 31 May 2016. Published online: 19 June 2016.

Authors:

Bustari Hasan, Faculty of Fisheries and Marine Sciences, Universitas Riau, Jalan HR. Subrantas KM 12.5 Pekanbaru, Indonesia, e-mail: bustarih@yahoo.com

Iskandar Putra, Faculty of Fisheries and Marine Sciences, Universitas Riau, Jalan HR. Subrantas KM 12.5 Pekanbaru, Indonesia, e-mail: putra@gmail.com

Indra Suharman, Faculty of Fisheries and Marine Sciences, Universitas Riau, Jl. HR. Subrantas KM 12.5 Pekanbaru, Indonesia, e-mail: indra70s@yahoo.com

Dian Iriani, Faculty of Fisheries and Marine Sciences, Universitas Riau, Jalan HR. Subrantas KM 12.5 Pekanbaru, Indonesia, e-mail: diriani@yahoo.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hasan B., Putra I., Suharman I., Iriani D., 2016 Evaluation of salted trash fish as a protein source replacing fishmeal in the diet for river catfish (*Hemibagrus nemurus*). *AAFL Bioflux* 9(3):647-656.

Evaluation of salted trash fish as a protein source replacing fishmeal in the diet for river catfish (*Hemibagrus nemurus*)

ORIGINALITY REPORT

23%
SIMILARITY INDEX

15%
INTERNET SOURCES

21%
PUBLICATIONS

3%
STUDENT PAPERS

PRIMARY SOURCES

1 sutir.sut.ac.th:8080 Internet Source **1%**

2 moscow.sci-hub.se Internet Source **1%**

3 ppid.unri.ac.id Internet Source **1%**

4 coek.info Internet Source **1%**

5 Azarm, Hamid Mohammadi, and Sang-Min Lee. "Effects of partial substitution of dietary fish meal by fermented soybean meal on growth performance, amino acid and biochemical parameters of juvenile black sea bream *Acanthopagrus schlegelii*", *Aquaculture Research*, 2012. Publication **1%**

6 Berto, Ricardo da Silva, Gabriella do Vale Pereira, José Luiz Pedreira Mouriño, Maurício Laterça Martins, and Débora Machado **1%**

Fracalossi. "Yeast extract on growth, nutrient utilization and haemato-immunological responses of Nile tilapia", Aquaculture Research, 2015.

Publication

7 kmbase.medic.or.kr 1 %
Internet Source

8 sinta3.ristekdikti.go.id 1 %
Internet Source

9 avesis.atauni.edu.tr 1 %
Internet Source

10 Tjipto Leksono, S Mus, Z Annisa. "Antibacterial Potential of Chitosan Extracted from Shrimp () Carapace ", Journal of Physics: Conference Series, 2019 <1 %
Publication

11 Bauer, W.. "Substitution of fishmeal with microbial floc meal and soy protein concentrate in diets for the pacific white shrimp *Litopenaeus vannamei*", Aquaculture, 20120415 <1 %
Publication

12 Jingping Guo, Weihua Gao, Benyue Guo, Wei Xu, Wenbing Zhang, Kangsen Mai. " Using a selectively bred nongenetically modified soybean meal to replace fishmeal in practical <1 %

diets for the Pacific white shrimp ",
Aquaculture Nutrition, 2018

Publication

13 Zhou, F.. "Effects of dietary arginine and lysine levels on growth performance, nutrient utilization and tissue biochemical profile of black sea bream, *Acanthopagrus schlegelii*, fingerlings", Aquaculture, 20110901

Publication

14 www.mdpi.com <1 %
Internet Source

15 biblio.ugent.be <1 %
Internet Source

16 www.abah.bioflux.com.ro <1 %
Internet Source

17 es.scribd.com <1 %
Internet Source

18 macau.uni-kiel.de <1 %
Internet Source

19 www.ncbi.nlm.nih.gov <1 %
Internet Source

20 www.scribd.com <1 %
Internet Source

21 www.thaiscience.info <1 %
Internet Source

22 Fagbenro, O.. "Physical and nutritional properties of moist fermented fish silage pellets as a protein supplement for tilapia (*Oreochromis niloticus*)", *Animal Feed Science and Technology*, 19980331 <1 %

Publication

23 De Luca, G.. "Amino acid levels in some brain areas of inducible nitric oxide synthase knock out mouse (iNOS^{-/-}) before and after pentylenetetrazole kindling", *Pharmacology, Biochemistry and Behavior*, 200612 <1 %

Publication

24 Guan, B.. "Metabolism traits of 'all-fish' growth hormone transgenic common carp (*Cyprinus carpio* L.)", *Aquaculture*, 20081101 <1 %

Publication

25 Joachim W. Hertrampf, Felicitas Piedad-Pascual. "Handbook on Ingredients for Aquaculture Feeds", Springer Nature, 2000 <1 %

Publication

26 M.A Hossain, U Focken, Klaus Becker. "Evaluation of an unconventional legume seed, *Sesbania aculeata*, as a dietary protein source for common carp, *Cyprinus carpio* L.", *Aquaculture*, 2001 <1 %

Publication

- 27 Rui Ma, Huapeng Hou, Kangsen Mai, Anant S. Bharadwaj, Hong Cao, Fengjie Ji, Wenbing Zhang. "Comparative study on the effects of L-methionine or 2-hydroxy-4-(methylthio) butanoic acid as dietary methionine source on growth performance and anti-oxidative responses of turbot (*Psetta maxima*)", *Aquaculture*, 2013
Publication <1 %
-
- 28 Submitted to Universiti Kebangsaan Malaysia
Student Paper <1 %
-
- 29 digitalcommons.uri.edu
Internet Source <1 %
-
- 30 Farhat, , and Mukhtar A Khan. "Growth, feed conversion and body composition of fingerling stinging catfish *Heteropneustes fossilis* (Bloch) fed varying levels of dietary l - threonine", *Aquaculture Research*, 2016.
Publication <1 %
-
- 31 Submitted to Pukyong National University
Student Paper <1 %
-
- 32 Roghieh Safari, Seyed Hossein Hoseinifar, Mohammad Reza Imanpour, Ali Hajibegloo et al. "The effects of multi-enzyme and betaine on growth performance, body composition haemato-immunological parameters and

expression of growth-related genes in beluga (Huso huso)", Aquaculture, 2022

Publication

33

aquaticcommons.org

Internet Source

<1 %

34

L.J. Ramseyer. "Effects of dietary protein to metabolizable energy ratios and total protein concentrations on the performance of yellow perch *Perca flavescens*", Aquaculture Nutrition, 12/1998

Publication

<1 %

35

Xiaohong Tan, Zhenzhu Sun, Meng Zhou, Cuiyun Zou, Hongyan Kou, Sarath Babu Vijayaraman, Yanhua Huang, Heizhao Lin, Li Lin. "Effects of dietary hawthorn extracts supplementation on lipid metabolism, skin coloration and gut health of golden pompano (*Trachinotus ovatus*)", Aquaculture, 2020

Publication

<1 %

36

A M Balogun. "Use of macadamia presscake as a protein feedstuff in practical diets for tilapia, *Oreochromis niloticus* (L.)", Aquaculture Research, 6/1995

Publication

<1 %

37

Anthony C. Ostrowski, S. Divakaran. "The amino acid and fatty acid compositions of selected tissues of the dolphin fish

<1 %

(Coryphaena hippurus) and their nutritional implications", Aquaculture, 1989

Publication

38

Madage, S. S. K., W. U. D. Medis, and Y. Sultanbawa. "Fish Silage as Replacement of Fishmeal in Red Tilapia Feeds", Journal of Applied Aquaculture, 2015.

Publication

<1 %

39

Submitted to Universitas Airlangga

Student Paper

<1 %

40

Xin - Dang Zhang, Jian - Wei Zhang, Heng - Zhi Wang, Bei - Bei Lin et al. " Evaluation of soybean meal as alternative to fish meal in diet for juvenile Asian red - tailed catfish () ", Aquaculture Nutrition, 2019

Publication

<1 %

41

Yucong Lin. "Dietary l-lysine requirement of juvenile Chinese sucker, Myxocyprinus asiaticus", Aquaculture Research, 04/2012

Publication

<1 %

42

discovery.researcher.life

Internet Source

<1 %

43

erepo.lib.uwu.ac.lk

Internet Source

<1 %

44

peneliti.unri.ac.id

Internet Source

<1 %

45 Cho, S.H.. "Effect of feeding ratio on growth and body composition of juvenile olive flounder *Paralichthys olivaceus* fed extruded pellets during the summer season", *Aquaculture*, 20060120
Publication <1 %

46 Hanan Sobhy, Mohammed Abo Elmagd, Maha El-kholy, Enas Abo Elmagd, Eman Laz. "Hepatoprotective Effect of Jojoba Oil and Nigella Sativa Seeds in Rats Fed Diet Containing Aflatoxin", *Egyptian Journal of Chemistry and Environmental Health*, 2016
Publication <1 %

47 ejbio.imedpub.com
Internet Source <1 %

48 espace.library.uq.edu.au
Internet Source <1 %

49 etd.aau.edu.et
Internet Source <1 %

50 oasis.repo.nii.ac.jp
Internet Source <1 %

51 www.ag.arizona.edu
Internet Source <1 %

52 www.sciencepublishinggroup.com
Internet Source <1 %

53

Jamjun Pechsiri. "A comparative study of growth and feed utilization efficiency of sex-reversed diploid and triploid Nile tilapia, *Oreochromis niloticus* L.", *Aquaculture Research*, 1/2005

Publication

<1 %

54

Jin Choi, Md. Mostafizur Rahman, Sang-Min Lee. "Distillers Dried Grain as a Partial Replacement for Wheat Flour in the Diet of Juvenile Rockfish *Sebastes schlegeli*", *Fisheries and aquatic sciences*, 2014

Publication

<1 %

55

M.R. Hasan, M.S. Haq, P.M. Das, G. Mowlah. "Evaluation of poultry-feather meal as a dietary protein source for Indian major carp, *Labeo rohita* fry", *Aquaculture*, 1997

Publication

<1 %

56

Mohammad Bagher Khoshhava, Abdolmohammad Abedian Kenari, Mohammad Kazem Mirzakhani. " The effects of concurrent of citric acid and soybean - based diets on growth performance, body composition, haemobiochemical indices, digestibility and fatty acid profile in juvenile rainbow trout, ", *Aquaculture Nutrition*, 2021

Publication

<1 %

57

Nguyen, Ngoc Thi Bich, Liet Chim, Pierrette Lemaire, and Laurent Wantiez. "Feed intake,

<1 %

molt frequency, tissue growth, feed efficiency and energy budget during a molt cycle of mud crab juveniles, *Scylla serrata* (Forskål, 1775), fed on different practical diets with graded levels of soy protein concentrate as main source of protein", *Aquaculture*, 2014.

Publication

58

Peres, H.. "Effect of the dietary essential amino acid pattern on growth, feed utilization and nitrogen metabolism of European sea bass (*Dicentrarchus labrax*)", *Aquaculture*, 20070703

Publication

59

S M Al-Ogaily. "Effect of feeding different grain sources on the growth performance and body composition of tilapia, *Oreochromis niloticus* (L.)", *Aquaculture Research*, 7/1996

Publication

60

adoc.pub

Internet Source

61

brage.bibsys.no

Internet Source

62

dergipark.org.tr

Internet Source

63

repositorio.ufmg.br

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

sciendo.com

64

Internet Source

<1 %

65

zsp.com.pk

Internet Source

<1 %

66

B.C. PETERSON, N.J. BOOTH, B.B. MANNING.
"Replacement of fish meal in juvenile channel catfish, *Ictalurus punctatus*, diets using a yeast-derived protein source: the effects on weight gain, food conversion ratio, body composition and survival of catfish challenged with *Edwardsiella ictaluri*", *Aquaculture Nutrition*, 2012

Publication

<1 %

67

D. M. S. D. El-Saidy, Ebtahal E. M. Hussein.
"REPLACEMENT OF FISH MEAL BY JOJOBA MEAL (*SIMMONDSIA CHINENSIS*) IN THE DIETS FOR RED TILAPIA, *OREOCHROMIS MOSSAMBICUS* X *O. NILOTICUS* FRY",
Menoufia Journal of Animal Poultry and Fish Production, 2018

Publication

<1 %

68

Kang-Woong Kim. "Optimum Dietary Protein Levels and Protein to Energy Ratios in Olive Flounder *Paralichthys olivaceus*", *Journal of the World Aquaculture Society*, 04/03/2007

Publication

<1 %

69

S. ZHANG. "Meat and bone meal replacement in diets for juvenile gibel carp (*Carassius auratus gibelio*): effects on growth performance, phosphorus and nitrogen loading", *Aquaculture Nutrition*, 10/2006

Publication

<1 %

70

Ying Chen, Gladstone Sagada, Bingying Xu, Wei Chao et al. " Partial replacement of fishmeal with single - cell protein in the diet for juvenile black sea bream () ", *Aquaculture Research*, 2019

Publication

<1 %

71

"Food Processing By - Products and their Utilization", Wiley, 2017

Publication

<1 %

72

Amanat Ali, Nasser A. Al-Asgah. " Effect of feeding different carbohydrate to lipid ratios on the growth performance and body composition of nile tilapia () fingerlings ", *Animal Research*, 2001

Publication

<1 %

73

Ehoulaman Marie-Angèle N'Gbaramou, Ahou Rachel Koumi, Boua Célestin Atsé, Kablan Tano. "Dry Feeds Produced Based on Local Available Feedstuffs for African Catfish *Clarias Gariepinus* Juveniles Rearing in Earthen Ponds", *European Journal of Agriculture and Food Sciences*, 2020

<1 %

74

Jonas R. Miller, Guillaume P. Salze, Kevin R. Stuart, Mark A. Drawbridge, D. Allen Davis. "The effect of dietary carbohydrate to lipid ratios on growth performance in juvenile California yellowtail *Seriola dorsalis*", *Aquaculture*, 2022

Publication

<1 %

75

Sang-Min Lee. "Effects of Dietary Protein and Energy Levels on Growth and Body Composition of Juvenile Flounder *Paralichthys olivaceus*", *Journal of the World Aquaculture Society*, 04/03/2007

Publication

<1 %

76

Shusaku Takagi, Hisashi Murata, Takanobu Goto, Makoto Endo, Hirofumi Yamashita, Masaharu Ukawa. "Taurine is an essential nutrient for yellowtail *Seriola quinqueradiata* fed non-fish meal diets based on soy protein concentrate", *Aquaculture*, 2008

Publication

<1 %

77

W Afuang. "Comparative nutritional evaluation of raw, methanol extracted residues and methanol extracts of moringa (*Moringa oleifera* Lam.) leaves on growth performance and feed utilization in Nile tilapia (*Oreochromis niloticus* L.)", *Aquaculture Research*, 11/2003

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On