

Growth performance and survival rate of Asian swamp eel in biofloc systems with different stocking density

by Indra Suharman


Submission date: 07-Apr-2023 01:03AM (UTC+0700)

Submission ID: 2057754770

File name: AACL_bioflux_vol_15-2-_2022.pdf (522.36K)

Word count: 5062

Character count: 25902



4

Growth performance and survival rate of Asian swamp eel in biofloc systems with different stocking density

Iesje Lukistyowati, Usman M. Tang, Iskandar Putra, Muhammad Fauzi, Indra Suharman, Rusliadi Rusliadi, Rudi Alfinda, Andi Nurahmad, Irwan Effendi

27

Faculty of Fisheries and Marine Sciences, University of Riau, Pekanbaru, Riau, Indonesia.
Corresponding author: I. Effendi, helpingirwan@gmail.com

10

Abstract. The cultivation of Asian swamp eel (*Monopterus albus*) has been known for a long time in Asia and the biofloc system is widely applied in the cultivation of some fish species in this region. However, so far, this technology has not been developed for Asian swamp eel cultivation. This study aimed to determine the growth performance and survival rate of this eel reared in a biofloc system with different stocking densities. The work was conducted from April to June 2020, by using a completely randomized design. The animals were reared in 100 L containers under different densities: treatment A (10 fish tank⁻¹), B (20 fish tank⁻¹), C (30 fish tank⁻¹), and D (40 fish tank⁻¹). The animals were fed with earthworms twice a day, 5% of bodyweight. The results showed that stocking densities affected the absolute weight (Wg) and specific growth rate (SGR) of Asian swamp eel significantly ($p < 0.05$). Treatment C was the best treatment compared to others, with an average absolute weight gain of 9.32 ± 0.461 g and an average daily growth rate of $2.69 \pm 0.238\%$. The best survival rate was in treatment C ($93.33 \pm 0.19\%$), although it was not significantly different from other treatments. The hematological profile (total erythrocytes, hemoglobin, hematocrit and total leukocytes) were also affected by the treatments ($p < 0.05$). The stocking density of 30 fish tank⁻¹ (C) yielded the highest total erythrocytes (2.88×10^6 cells mm⁻³), hemoglobin (17.4 g dL⁻¹), hematocrit (31.66%), and total leukocytes (3.57×10^4 cells mm⁻³). The difference in stocking density did not have an effect on blood glucose levels ($p > 0.05$). Glucose levels ranged from 33.33 to 54.66 mg dL⁻¹, with the highest glucose levels in treatment D.

Key Words: absolute weight, blood profiles, daily growth rate, hematological profiles, ricefield eel.

15

Introduction. The Asian swamp eel (*Monopterus albus*) is a commercially important, air-breathing species of fish from the family Synbranchidae. The animal is also known as rice eel, rice field eel, or rice paddy eel. The native range includes tropical, subtropical and temperate climates, but it is mainly found in East and Southeast Asia, where it is a very common fish sold throughout the region (Matsumoto et al 2010; Chen et al 2021). Eels are a good protein source, with a protein content of 66.7%, fat content of 10.74%, and an ash content of 7% (Vishwanath et al 1998).

The demand for eels increased through the years, and it is estimated that only 10% can be provided (Suzuki et al 2003). The eel cultivation is still using mud media with a mixture of coarse and fine bran, straw, banana stems, and manure as a substrate (Herawati et al 2018). The stocking density is one of the factors that determines the growth of fish in a rearing system. Determination of the proper stocking density will not only provide information on optimal growth for fish, but will also increase business efficiency. Such information, especially for the Asian swamp eel, is still limited.

Biofloc is a particle that is stirred by aeration and circulation, which consists of a collection of autotrophic and heterotrophic organisms (in the form of bacteria, phytoplankton, fungi, ciliates, nematodes and detritus) and non-living materials (Avnimelech et al 2015). This management system converts organic and inorganic compounds containing carbon compounds (C), hydrogen (H), oxygen (O), nitrogen (N) with little available phosphorus (P) into sludge mass in the form of bioflocs by using

1

flocs-forming bacteria, which synthesize poly hydroxy alkanoate biopolymers, namely as bioflocs bonds (Abakari et al 2021).

Biofloc in fish culture is applied by adding a starter in the form of carbon and bacteria. The addition of organic carbohydrates into the medium could stimulate bacterial growth (de Lima et al 2019; Hargreaves 2006; Rajkumar et al 2015). Biofloc technology provides natural feed for domestic fish, is feasible, and does not pollute the environment (Emerenciano et al 2017; Supriatna et al 2014). The application of biofloc technology has been widely used in fish and shrimp culture (Ekasari et al 2015; Putra et al 2017; Serra et al 2015), but the information in application of the technology to Asian swamp eel is still limited. This study aimed to determine the growth performance and survival rate of this eel reared in biofloc system under different stocking densities.

Material and Method

Time and place. The research was carried out from April to June 2020, in the Aquaculture Technology Laboratory, Faculty of Fisheries and Marine Sciences, Riau University, Indonesia. The animals were purchased from local an Asian swamp eel hatchery in Riau.

Experimental design. The experiment was conducted within the ethical guidelines in animal research developed by the National Centre for the Replacement, Refinement and Reduction of Animals in Research (NC3Rs). A completely randomized design with four levels of treatment and three replications was used in this study. The Asian swamp eels were reared in 100 L container under different densities, namely treatment A (10 fish tank⁻¹), B (20 fish tank⁻¹), C (30 fish tank⁻¹), and D (40 fish tank⁻¹).

Biofloc cultivation system. The biofloc cultivation systems were set up in a 100 L canvas containers. Each container was filled with 25 L of water. The system was prepared by mixing 10 mL of probiotics *Bacillus* sp. (5x10¹⁰ CFU mL⁻¹) and 0.48 mL L⁻¹ water molasses into the containers and aerated continuously for seven days to grow the flock. The Asian swamp eels (average weight of 5.85±0.75 g) were stocked into the system, according to the above treatments. Additional inoculants, namely 10 mL of probiotics *Bacillus* sp. (5x10¹⁰ CFU mL⁻¹) were added to each tank at every 5-day interval. The animals were fed with earthworms (*Lumbricus rubellus*) twice a day, 5% of total body weight. Fish weight gain was measured every ten days for 40 days.

Growth of Asian swamp eel. Growth performance of Asian swamp eel was measured by weighing the animals at the beginning and at the end of experiment. The weight gain (Wg) was calculated as follows:

$$Wg = Wt - Wo$$

Where: Wg - weight gain (g); Wt - the weight of the eel at the end of the experiment (g); and Wo - the weight of the eel at the beginning of the experiment (g).

Specific growth rate (SGR) was calculated by using the following formula;

$$SGR (\%/day) = \frac{(\ln Wt - \ln Wo)}{t} \times 100 \%$$

Where: Wt - the weight of the eel at the end of the experiment (g); Wo - the weight of eel at the beginning of the experiment (g); and t - time or duration of cultivation (days).

Feed efficiency. The efficiency of feed utilization consumed by the Asian swamp eel was measured referring to de Verdal et al (2017):

$$FE = \frac{((Wt+D)-Wo)}{F} \times 100$$

Where: FE - feed efficiency (%); ⁵ Wt - weight of the fish biomass at the end of the experiment (g); Wo - weight of the fish biomass at the beginning of the experiment (g); D - weight of the fish that died during the study (g); F - amount of feed provided to the fish during the study (g).

⁴³ **Hematological profile.** Blood samples were collected from 3 animals per container at the end of the study (40 days). The Asian swamp eel was anesthetized with 0.1 mL L⁻¹ clove oil before blood collection. A set of clean and sterile syringe and Eppendorf ³⁷ e were rinsed with blood anti-coagulant 10% EDTA. Blood sample (as much as 1 mL) was collected with a 1 mL syringe from the caudal vein at a slope of 45° to pierce the vein. A volume of 0.5 mL of blood was placed into an Eppendorf tube. Hematological profiles of blood included total erythrocyte cells mm⁻³ (Blaxhall 1972), hemoglobin (%) (Wedemeyer & Yasutake 1977), hematocrit levels (%) (Mauel et ⁴⁸ 2007), total leukocytes (cells mm⁻³) (Blaxhall 1972), leukocytes and blood glucose (mg dL⁻¹) (Papazafiropoulou et al 2010).

¹⁸ **Water quality.** The main water quality parameters such as dissolved oxygen (DO), pH, and temperature were measured us⁴² a digital water checker (YSI-550 A, ASTM, Alla, France) at 10-day intervals, while total ammonia nitrogen (TAN) was measured every ten days using spectrophotometric method (APHA 2008).

⁴⁴ **Total ammonia nitrogen.** Total ammonia nitrogen (TAN) was measured every 10 days for each treatment for 40 days. It was measured using a spectrophotometer (Optima SP300), with a wavelength of 630 nm. The measurements were carried out by placing 50 mL of the sample into a 100 mL Erlenmeyer flask, then adding 1 mL of Nessler's solution, then shaking it and allowing the reaction process to last for at least 10 min. The solution was put into a cuvette on a spectrophotometer at a wavelength of 630 nm, read and recorded on the absorption scale. The levels of TAN were calculated using a calibration curve or a straight line equation (APHA 2008) and determined by the formula:

$$\text{TAN concentration (mg L}^{-1}\text{)} = (\text{Cst} \times \text{As})/\text{Ast}$$

Where: Cst - concentration of the standard solution; Ast - absorbance value of standard solution; As - the absorbance value of the sample solution.

²⁴ **Data analysis.** The data were subject to one-way analysis of variance (ANOVA) test to determine the effect ⁸ treatments on some parameters and followed by Newman-Keuls multiple range test, with a confidence level of 95%, while the water quality data was analyzed descriptively.

Results and Discussion

² **Growth and survival rates of Asian swamp eel.** The results showed that the Wg of Asian swamp eel ranged from 6.96±0.695 to 9.32±0.461 g, SGR ranged from 1.9±0.379 to 2.69±0.238%, and survival rates ranged ¹⁰ from 90 to 93.33±0.19%. ANOVA results showed that weight gain and SGR were significantly different between treatments (p<0.05), while survival rate was not significantly different (p>0.05). Treatment C (30 fish tank⁻¹) was the best one compared to others, with an average absolute weight of 9.32±0.461 g and an average ¹³ SGR of 2.69±0.238%. The survival rate in treatment C was 93.33±0.19%, although it was not significantly different from other treatments. The best feed efficiency was also observed in treatment C (Table 1).

Hematological profiles. The cultivation of Asian swamp eel with different stocking densities had an effect on the hematological profile (total erythrocytes, hemoglobin, hematocrit and total leukocytes) (p<0.05). The stocking density of 30 individuals tank⁻¹ (C) yielded the highest total erythrocytes (2.88 x 10⁶ cells mm⁻³), hemoglobin (17.4 g dL⁻¹), hematocrit (31.66%), and total leukocytes (3.57 x 10⁴ cells mm⁻³). The hematological profile of the eels during the study was in the normal range. Siang et al (2007) reported that the normal hematological profiles of *Monopterus albus* range as follows: erythrocytes between 1.05 and 1.10 x 10⁶ cell mm⁻³, leukocyte between 47.58

and 48.56×10^3 cell mm^{-3}), hemoglobin between 20 and 21 g dL^{-1}), and hematocrit between 47.05 and 48.10 %. The difference in stocking density did not have an effect on blood glucose levels ($p > 0.05$). Glucose levels ranged from 33.33 to 54.66 mg dL^{-1} , the highest glucose levels occurring in treatment D (Table 2).

Table 1
Growth performance and survival rate of Asian swamp eel

No	Parameter	Treatment			
		A	B	C	D
1	Weight gain (g)	7.38 ± 0.500^a	8.32 ± 0.429^b	9.32 ± 0.461^c	6.96 ± 0.695^a
2	SGR (%)	1.90 ± 0.379^a	2.24 ± 0.105^a	2.69 ± 0.238^b	1.88 ± 0.352^a
3	Survival rate (%)	90.00 ± 0.00^a	90.00 ± 0.00^a	93.33 ± 0.19^a	92.50 ± 0.00^a
4	Feed efficiency (%)	42.41 ± 3.799^a	49.67 ± 2.105^a	67.46 ± 9.806^b	41.00 ± 5.949^a

Note: SGR - specific growth rate; data represent mean and standard deviation; different superscripts in the same row show significant differences ($p < 0.05$).

Table 2
Hematological profiles of Asian swamp eel

No	Parameter	Treatment			
		A	B	C	D
1	Erythrocyte (cell mm^{-3})	$181.66 \times 10^4 \pm 1.53^b$	$271.66 \times 10^4 \pm 2.08^c$	$288.33 \times 10^4 \pm 1.53^d$	$120.33 \times 10^4 \pm 1.53^a$
2	Hemoglobin (g dL^{-1})	13.46 ± 1.28^a	14.8 ± 0.72^a	17.4 ± 1.44^b	13.4 ± 0.52^a
3	Hematocrit (%)	25.33 ± 1.52^b	27 ± 2^b	31.66 ± 1.52^c	20.33 ± 0.57^a
4	Leukocyte	3.27 ± 0.02^a	3.38 ± 0.08^a	3.57 ± 0.07^b	3.62 ± 0.12^b
5	Glucose (mg dL^{-1})	33.33 ± 4.16^a	41.33 ± 1.52^a	46.23 ± 2.51^a	54.66 ± 3.05^a

Note: data represent mean and standard deviation; different superscripts in the same row show significant differences ($p < 0.05$).

Water quality. In this study, water quality parameters were in normal ranges for aquatic organism life. The temperature ranged from $26.64 \pm 0.3^\circ\text{C}$ to $26.95 \pm 0.32^\circ\text{C}$, pH ranged from 7.39 ± 0.08 to 7.60 ± 0.15 , dissolved oxygen ranged from $7 \pm 0.57 \text{ mg L}^{-1}$ to $7.53 \pm 0.21 \text{ mg L}^{-1}$ and ammonia values from $0.231 \pm 0.012 \text{ mg L}^{-1}$ to $0.003 \pm 0.018 \text{ mg L}^{-1}$. The results of the analysis on water quality were not significantly different ($p < 0.05$) among the experimental treatments. The average values of the water quality parameters are presented in Table 3.

Table 3
Water quality of culture media of the biofloc system

No	Parameter	Treatment			
		A	B	C	D
1	Temperature ($^\circ\text{C}$)	26.78 ± 0.58	26.95 ± 0.51	26.64 ± 0.3	26.84 ± 0.35
2	pH	7.39 ± 0.08	7.45 ± 0.09	7.6 ± 0.15	7.45 ± 0.04
3	Dissolved oxygen (mg L^{-1})	7.53 ± 0.21	7.47 ± 0.31	7 ± 0.57	7.2 ± 0.46
4	Ammonia (mg L^{-1})	0.001 ± 0.012	0.002 ± 0.018	0.003 ± 0.017	0.003 ± 0.018

The results showed that the best growth, highest survival and best level of feed efficiency in Asian swamp eels was in treatment C (the stocking density of 30 fish tank^{-1}). The increase of stock density to 40 fish tank^{-1} inhibited or slowed down the growth rate. It is thought that increased density was associated with stress levels, food competition, space availability for movement, appetite, and antagonistic interactions (Breine et al 1996; Ronald et al 2014; Yousif 2017). Stocking density does affect fish growth. Hastuti & Subandiyono (2020) reported that density of tilapia cultivated in an aquaponic system had an effect on biological parameters, such as growth, feed consumption and feed utilization efficiency. Billah et al (2019) also noted that the suitable stocking density was 6 m^{-2} for *C. carpio* and *O. niloticus* in rice fields for better growth and survival. However, stocking density does not affect the survival rates and hematological performances. This

has been reported by several researchers; for example, El-Dahhar et al (2021) reported that growth performance indices of meagre (*Argyrosomus regius*) decreased with increasing fish density, but without significant difference ($p>0.05$). The growth of eels can be influenced by several factors, including feed quality, water quality, stocking density, and the application of biofloc used in the culture system (Hadiroseyani et al 2016; Putra et al 2020). The density of organisms maintained in the culture system also affected the growth, production, and space efficiency. Some authors reported that low stocking density may also cause a reduced growth (Ronald et al 2014).

Application of biofloc technology had a significant influence on the growth and survival of eels, since this technology is able to increase water quality parameters (Nurhatijah et al 2016; Stein & Klotz 2016), and can provide natural feed as additional feed, even though eels are fed with sufficient food (Avnimelech 1999; Putra et al 2017). The feed provided also plays a main role in the growth of eels. The feed used was earthworms with a high nutrient content, meeting the nutritional needs for Asian swamp eel. The dry matter of earthworms contains nutrients such as protein 60-72%, 7-10% fat, and 8-10% ash, making it suitable for carnivorous eels, including carnivorous Asian swamp eels that require high protein feed for growth and development (Hill & Watson 2007).

The stocking densities had an effect on the hematological profile (total erythrocytes, hemoglobin, hematocrit and total leukocytes) ($p<0.05$). The stocking density of 30 individuals per tank (C) provided the highest yield for total erythrocytes, hemoglobin, hematocrit, and total leukocytes. This data corresponds to the absolute weight gain and survival rate data of the Asian swamp eels. These animals grow better in these conditions and this can be seen in through their hematological profile.

The stocking density can be a trigger that affects movement and physiological activity, thereby affecting the blood profile. Some factors (species, sex, dietary nutrition, size, physical activity, and age of the fish) may influence the number of erythrocytes, hemoglobin, and hematocrit level. Overall, physiological status and environmental conditions do affect the number of erythrocytes (Fazio 2019; Syawal et al 2021).

The difference in stocking density in Asian swamp eel cultivation among the experimental treatments did not have an effect on blood glucose levels ($p>0.05$). Glucose levels ranged from 33.33-54.66 mg dL⁻¹, in which the highest glucose levels were in treatment D. Glucose levels in this study were lower compared to the findings of Hu et al (2018), where the levels ranged from 54 to 144 mg dL⁻¹. Peng et al (2019) reported that normal Asian swamp eel glucose levels range from 45 to 68 mg dL⁻¹, while Shi et al (2019) revealed that glucose levels ranged from 43 to 51 mg dL⁻¹. It can be said that Asian swamp eel glucose levels in treatment B, C, and D were within the normal range.

Water quality parameters also affect the growth of fish. Fish culture with biofloc technology provides several advantages, including improved water quality and additional feed (Stein & Klotz 2016; Putra et al 2017). The quality of rearing water for eels is in the optimal range for growth. Biofloc contributed to the growth and production of the cultured organism, as shown in this study. This technology's basic principle is using heterotrophic bacteria to manage the C:N ratio in the water media (Avnimelech 1999; Hargreaves 2006). Therefore, biofloc technology is considered suitable for the Asian swamp eel cultivation.

Conclusions. Cultivation in biofloc system with different stocking densities affected the weight gain and specific growth rate of Asian swamp eel significantly ($p<0.05$). Treatment C (30 fish tank⁻¹) was the best treatment compared to others, with an average absolute weight of eels of 9.32±0.461 g and an average specific growth rate of 2.69±0.238%. The survival rate in treatment C was 93.33±0.19 %, although it was not significantly different from other treatments. The hematological profile (total erythrocytes, hemoglobin, hematocrit and total leukocytes) revealed that the animals had a significantly better immunity status ($p<0.05$). The stocking density of 30 individuals tank⁻¹ (C) provided the highest total erythrocytes (2.88 x10⁶ cells mm⁻³), hemoglobin (17.4 g dL⁻¹), hematocrit (31.66%), and total leukocytes (3.57 x 10⁴ cells mm⁻³).

Stocking density did not affect the blood glucose levels of these animals ($p > 0.05$). Glucose levels ranged from 33.33-54.66 mg dL⁻¹, with the highest glucose levels in treatment D.

30

Acknowledgements. The authors would like to thank the Institutes of Research and Public Services, University of Riau, for funding this study. We also express our gratitude to the laboratory technicians and students who have helped during the implementation of this research.

Conflict of Interest. The authors declare that there is no conflict of interest.

References

- Abakari G., Luo G., Kombat E. O., 2021 Dynamics of nitrogenous compounds and their control in biofloc technology (BFT) systems: A review. *Aquaculture and Fisheries* 6(5):441-447.
- Avnimelech Y., 1999 Carbon/nitrogen ratio as a control element in aquaculture systems. *Aquaculture* 176(3-4):227-235.
- Avnimelech Y., De-Schryver P., Emmereciano M., Kuhn D., Ray A., Taw N., 2015 Biofloc technology - A practical guidebook. 3rd Edition. World Aquaculture Society, Baton Rouge, 259 p.
- Billah M. M., Uddin M. K., Samad M. Y. A., Hassan M. Z. B., Anwar M. P., Kamal A. H. M., Shahjahan M., Asif A. A., 2019 Effects of different stocking density of Nile tilapia (*Oreochromis niloticus*) and common carp (*Cyprinus carpio*) on the growth performance and rice yield in rice-fish farming system. *AAFL Bioflux* 13(2):789-803.
- Blaxhall P. C., 1972 The haematological assessment of the health of freshwater fish. A review of selected literature. *Journal of Fish Biology* 4(4):593-604.
- Breine J. J., Nguenga D., Teugels G. G., Ollevier F. A., 1996 Comparative study on the effect of stocking density and feeding regime on the growth rate of *Tilapia cameronensis* and *Oreochromis niloticus* (Cichlidae) in fishculture in Cameroon. *Aquatic Living Resources* 9(1):51-56.
- Chen H., Liu H., Li R., Lin X., Luo H., Ji H., Hu W., Luo D., 2021 Blood cell identification and hematological analysis during natural sex reversal in rice field eel (*Monopterus albus*). *Aquaculture* 538:736543.
- de Lima P. C. M., Silva L. O. B., de Lima A. J., da Silva S. M. B. C., Severi W., Gálvez A. O., 2019 Tilapia cultivated in a low-salinity biofloc system supplemented with *Chlorella vulgaris* and different molasses application rates. *Boletim do Instituto de Pesca* 45(4):e494, 11 p.
- De Verdal H., Komen H., Quillet E., Chatain B., Allal F., Benzie J. A. H., Vandeputte M., 2017 Improving feed efficiency in fish using selective breeding: a review. *Reviews in Aquaculture* 10(4):833-851.
- Ekasari J., Zairin Jr. M., Putri D. U., Sari N. P., Surawidjaja E. H., Bossier P., 2015 Biofloc-based reproductive performance of Nile tilapia *Oreochromis niloticus* L. broodstock. *Aquaculture Research* 46(2):509-512.
- El-Dahhar A. A. E., El-Ebiary E. H., Abdel-Rahim M. M., Refaee W. M. A., Lotfy A. M. A., 2021 Stocking density effects on survival, growth performance, feed utilization, and carcass composition of meagre, *Argyrosomus regius* (Asso, 1801) fingerlings reared in fiberglass tanks using underground saltwater. *AAFL Bioflux* 14(1):495-505.
- Emerenciano M. G. C., Martínez-Córdova L. R., Martínez-Porchas M., Miranda-Baeza A., 2017 Biofloc technology (BFT): a tool for water quality management in aquaculture. *Water Quality* 5:92-109.
- Fazio F., 2019 Fish hematology analysis as an important tool of aquaculture: A review. *Aquaculture* 500:237-242.
- Hadiroseyani Y., Sukenda S., Surawidjaja E. H., Utomo N. B. P., Affandi R., 2016 Survival rate of transported ricefield eels, *Monopterus albus* (Synbranchidae), in open and closed system at water salinity level of 0 and 9 g L⁻¹. *AAFL Bioflux* 9(3):759-767.

- Hargreaves J. A., 2006 Photosynthetic suspended-growth systems in aquaculture. *Aquacultural Engineering* 34(3):344-363.
- Hastuti S., Subandiyono S., 2020 Aminotransferase, hematological indices and growth of tilapia (*Oreochromis niloticus*) reared in various stocking densities in aquaponic systems. *AACL Bioflux* 13(2):813-824.
- Herawati V. E., Nugroho R. A., Pinandoyo, Hutabarat J., Prayitno B., Karnaradjasa O., 2018 The growth performance and nutrient quality of Asian swamp eel *Monopterus albus* in Central Java Indonesia in a freshwater aquaculture system with different feeds. *Journal of Aquatic Food Product Technology* 27(6):658-666.
- Hill J. E., Watson C. A., 2007 Diet of the nonindigenous Asian swamp eel in tropical ornamental aquaculture ponds in west-central Florida. *North American Journal of Aquaculture* 69(2):139-146.
- Hu Y., Yang G., Li Z., Hu Y., Zhong L., Zhou Q., 2018 Effect of dietary taurine supplementation on growth, digestive enzyme, immunity and resistant to dry stress of rice field eel (*Monopterus albus*) fed low fish meal diets. *Aquaculture Research* 49(5):2108-2118.
- Matsumoto S., Kon T., Yamaguchi M., Takeshima H., Yamazaki Y., Mukai T., Kuriwa K., Kohda M., Nishida M., 2010 Cryptic diversification of the swamp eel *Monopterus albus* in East and Southeast Asia, with special reference to the Ryukyuan populations. *Ichthyological Research* 57(1):71-77.
- Mauel M. J., Miller D. L., Merrill A. L., 2007 Hematologic and plasma biochemical values of healthy hybrid tilapia (*Oreochromis aureus* x *Oreochromis nilotica*) maintained in a recirculating system. *Journal of Zoo and Wildlife Medicine* 38(3):420-424.
- Nurhatijah N., Muchlisin Z. A., Sarong M. A., Supriatna A., 2016 Application of biofloc to maintain the water quality in culture system of the tiger prawn (*Penaeus monodon*). *AACL Bioflux* 9(4):923-928.
- Papazafiropoulou A., Kardara M., Sotiropoulos A., Bousboulas S., Stamataki P., Pappas S., 2010 Plasma glucose levels and white blood cell count are related with ankle brachial index in type 2 diabetic subjects. *Hellenic Journal of Cardiology* 51(5):402-406.
- Peng M., Xue J., Hu Y., Wen C., Hu B., Jian S., Yang G., 2019 Disturbance in the homeostasis of intestinal microbiota by a high-fat diet in the rice field eel (*Monopterus albus*). *Aquaculture* 502:347-355.
- Putra I., Rusliadi R., Fauzi M., Tang U. M., Muchlisin Z. A., 2017 Growth performance and feed utilization of African catfish *Clarias gariepinus* fed a commercial diet and reared in the biofloc system enhanced with probiotic. *F1000Research* 6:1545, 9 p.
- Putra P., Effendi I., Lukistyowati I., Tang U. M., Fauzi M., Suharman I., Muchlisin Z. A., 2020 Effect of different biofloc starters on ammonia, nitrate, and nitrite concentrations in the cultured tilapia *Oreochromis niloticus* system. *F1000Research* 9:293.
- Rajkumar M., Pandey P. K., Aravind R., Vennila A., Bharti V., Purushothaman C. S., 2015 Effect of different biofloc system on water quality, biofloc composition and growth performance in *Litopenaeus vannamei* (Boone, 1931). *Aquaculture Research* 47(11):3432-3444.
- Ronald N., Gladys B., Gasper E., 2014 The effects of stocking density on the growth and survival of Nile tilapia (*Oreochromis niloticus*) fry at Son Fish Farm. *Journal of Aquaculture Research & Development* 5:2, 7 p.
- Serra F. P., Gaona C. A. P., Furtado P. S., Poersch L. H., Wasielesky W., 2015 Use of different carbon sources for the biofloc system adopted during the nursery and grow-out culture of *Litopenaeus vannamei*. *Aquaculture International* 23(6):1325-1339.
- Shi Y., Zhong L., Ma X., Liu Y., Tang T., Hu Y., 2019 Effect of replacing fishmeal with stickwater hydrolysate on the growth, serum biochemical indexes, immune indexes, intestinal histology and microbiota of rice field eel (*Monopterus albus*). *Aquaculture Reports* 15:100223.
- Siang H. Y., Yee L. M., Seng C. T., 2007 Acute toxicity of organochlorine insecticide endosulfan and its effect on behaviour and some hematological parameters of Asian

- swamp eel (*Monopterus albus*, Zuiew). *Pesticide Biochemistry and Physiology* 89:46-53.
- Stein L. Y., Klotz M. G., 2016 The nitrogen cycle. *Current Biology* 26(3):R94-R98.
- Supriatna A., Nurhatijah N., Sarong M. A., Muchlisin Z. A., 2019 Effect of biofloc density and crude protein level in the diet on the growth performance, survival rate, and feed conversion ratio of black tiger prawn (*Penaeus monodon*). *IOP Conference Series: Earth and Environmental Science* 348:12131, 7 p.
- Suzuki Y., Maruyama T., Numata H., Sato H., Asakawa M., 2003 Performance of a closed recirculating system with foam separation, nitrification and denitrification units for intensive culture of eel: towards zero emission. *Aquacultural Engineering* 29(3-4):165-182.
- Syawal H., Kurniawan R., Effendi I., Austin B., 2021 Fermented medicinal herbs improve the hematological and physiological profile of striped catfish (*Pangasionodon hypophthalmus*). *F1000Research* 10:466.
- Vishwanath W., Lilabati H., Bijen M., 1998 Biochemical, nutritional and microbiological quality of fresh and smoked mud eel fish *Monopterus albus* - a comparative study. *Food Chemistry* 61(1-2):153-156.
- Wedemeyer G. A., Yasutake W. T., 1977 Clinical methods for the assessment on the effect of environmental stress on fish health. Department of the Interior, Fish and Wildlife Service, 18 p.
- Yousif O. M., 2017 The effects of stocking density, water exchange rate, feeding frequency and grading on size hierarchy development in juvenile Nile tilapia, *Oreochromis niloticus* L. *Emirates Journal of Food and Agriculture* 4(1):45-53.
- *** APHA (American Public Health Association), 2008 Standard methods for the examination of water and wastewater. Washington D.C.

Received: 14 July 2021. Accepted: 25 September 2021. Published online: 25 April 2022.

Authors:

Iesje Lukistyowati, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: iesje.lukistiowati@lecturer.unri.ac.id
Usman Muhammad Tang, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: usman_mt@yahoo.co.id
Iskandar Putra, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: iskandar.putra@lecturer.unri.ac.id
Muhammad Fauzi, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: m.fauzi@lecturer.unri.ac.id
Indra Suharman University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: indra.suharman@lecturer.unri.ac.id
Rusliadi Rusliadi, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: rusliadi@lecturer.unri.ac.id
Rudi Alfinda, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: rudi.alfinda@student.unri.ac.id
Andi Nurahmad, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: andiinurr178@gmail.com
Irwan Effendi, University of Riau, Faculty of Fisheries and Marine Sciences, 28292 Pekanbaru, Indonesia, e-mail: helpingirwan@gmail.com.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Lukistyowati I., Tang U. M., Putra I., Fauzi M., Suharman I., Rusliadi R., Alfinda R., Nurahmad A., Effendi I., 2022 Growth performance and survival rate of Asian swamp eel in biofloc systems with different stocking density. *AAFL Bioflux* 15(2):1013-1020.

Growth performance and survival rate of Asian swamp eel in biofloc systems with different stocking density

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

15 %
INTERNET SOURCES

15 %
PUBLICATIONS

8 %
STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Universitas Jenderal Soedirman
Student Paper **2** %
- 2** Vivi Endar Herawati, Ristiawan Agung Nugroho, Pinandoyo, Johannes Hutabarat, Budi Prayitno, Ocky Karnaradjasa. " The Growth Performance and Nutrient Quality of Asian Swamp Eel in Central Java Indonesia in a Freshwater Aquaculture System with Different Feeds ", Journal of Aquatic Food Product Technology, 2018
Publication **1** %
- 3** Submitted to Syiah Kuala University
Student Paper **1** %
- 4** www.mdpi.com
Internet Source **1** %
- 5** www.worldnewsnaturalsciences.com
Internet Source **1** %
- 6** www.ijsr.net
Internet Source **1** %

7	www2.stat.unibo.it Internet Source	1 %
8	RC Mukti, M Amin, Yulisman. "Utilization of alternative ingredients as combination feed on the growth of catfish (<i>Clarias</i> sp.)", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
9	Submitted to Universitas Mulawarman Student Paper	1 %
10	journal.ipb.ac.id Internet Source	1 %
11	edepot.wur.nl Internet Source	1 %
12	www.cellphysiolbiochem.com Internet Source	1 %
13	pt.scribd.com Internet Source	<1 %
14	Tran Nguyen Duy Khoa, Chau Tai Tao, Ly Van Khanh, Tran Ngoc Hai. "Super-intensive culture of white leg shrimp (<i>Litopenaeus vannamei</i>) in outdoor biofloc systems with different sunlight exposure levels: Emphasis on commercial applications", Aquaculture, 2020 Publication	<1 %

15	alchetron.com Internet Source	<1 %
16	ijmcmed.org Internet Source	<1 %
17	I Putra, I Efendi, I Lukisyowati, U M Tang, R Rusliadi, R A Manik. " Effects of Various Carbon Doses on Tilapia (sp) Culture with Biofloc Technology ", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %
18	www.tceq.texas.gov Internet Source	<1 %
19	Submitted to Asian Institute of Technology Student Paper	<1 %
20	Submitted to Curtin University of Technology Student Paper	<1 %
21	Juliana Heloisa Pinê Américo-Pinheiro, Claudinei da Cruz, Mario Mamede Aguiar, Nádia Hortense Torres et al. "Sublethal Effects of Imidacloprid in Hematological Parameters of Tilapia (Oreochromis niloticus)", Water, Air, & Soil Pollution, 2019 Publication	<1 %
22	Neda Mirzakhani, Eisa Ebrahimi, Seyed Amir Hossein Jalali, Julie Ekasari. "Growth performance, intestinal morphology and	<1 %

nonspecific immunity response of Nile tilapia (Oreochromis niloticus) fry cultured in biofloc systems with different carbon sources and input C:N ratios", Aquaculture, 2019

Publication

23

S. Kumari, V. Harikrishna, V.K.R. Surasani, A.K. Balange, A.M. Babitha Rani. "Growth, biochemical indices and carcass quality of red tilapia reared in zero water discharge based biofloc system in various salinities using inland saline ground water", Aquaculture, 2021

Publication

24

Zhe Yu, Liang Li, Rui Zhu, Min Li, Li - Fang Wu. " Effects of bioflocs with different C/N ratios on growth, immunological parameters, antioxidants and culture water quality in Dybowski ", Aquaculture Research, 2019

Publication

25

aquicultura.furg.br

Internet Source

26

biblio.ugent.be

Internet Source

27

mulyadi.staff.unri.ac.id

Internet Source

28

D.P Hartono, E Barades. "Effectiveness of using Commercial Probiotics in Biofloc System

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

Culture Media on Growth, FCR, and Feed Efficiency of Catfish (*Clarias Gariepinus*)", IOP Conference Series: Earth and Environmental Science, 2022

Publication

29

Siti Aslamyah, Zainuddin, Badraeni, Muhammad Khairul Umam, Ayu Dewi Lestari, Haslinda. "The effect of protein levels in fermentation feed supplemented *lumbricus* sp. extract as feed additive on growth performance and body chemical composition of milkfish, *Chanos chanos forskal 1775*", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

30

journals.sagepub.com

Internet Source

<1 %

31

www.nature.com

Internet Source

<1 %

32

Joya Saha, M.A. Hossain, M. Al. Mamun, M.R. Islam, M.S. Alam. "Effects of carbon-nitrogen ratio manipulation on the growth performance, body composition and immunity of stinging catfish *Heteropneustes fossilis* in a biofloc-based culture system", *Aquaculture Reports*, 2022

Publication

<1 %

33

Mohammad Hossein Khanjani, Moslem Sharifinia. "Biofloc technology as a promising tool to improve aquaculture production", *Reviews in Aquaculture*, 2020

Publication

<1 %

34

Submitted to St. Ignatius College Preparatory

Student Paper

<1 %

35

iskandar.staff.unri.ac.id

Internet Source

<1 %

36

www.fal.infish.com.pl

Internet Source

<1 %

37

C.-Y. Huang. "Effects on acid-base balance, methaemoglobinemia and nitrogen excretion of European eel after exposure to elevated ambient nitrite", *Journal of Fish Biology*, 9/2002

Publication

<1 %

38

Carlos A. Mendez, María Cristina Morales, Germán E. Merino. "SETTLING VELOCITY DISTRIBUTION OF BIOFLOCULES GENERATED WITH DIFFERENT CARBON SOURCES DURING THE REARING OF THE RIVER SHRIMP *Cryphiops caementarius* WITH BIOFLOC TECHNOLOGY", *Aquacultural Engineering*, 2021

Publication

<1 %

39

Maryam Irani, Houman Rajabi Islami, Mahmoud Nafisi Bahabadi, Seyed Pezhman Hosseini Shekarabi. "Production of Pacific white shrimp under different stocking density in a zero-water exchange biofloc system: effects on water quality, zootechnical performance, and body composition", *Aquacultural Engineering*, 2022

Publication

<1 %

40

Oscar Satio Cunha Lobato, Felipe de Azevedo Silva Ribeiro, Anselmo Miranda-Baeza, Mauricio Gustavo Coelho Emerenciano. "Production performance of *Litopenaeus vannamei* (Boone, 1931) fed with different dietary levels of tilapia processing waste silage reared in biofloc system using two carbon sources", *Aquaculture*, 2019

Publication

<1 %

41

Roghayeh Dehghani, Amin Oujifard, Mansour Torfi Mozanzadeh, Vahid Morshedi, Dara Bagheri. "Effects of dietary taurine on growth performance, antioxidant status, digestive enzymes activities and skin mucosal immune responses in yellowfin seabream, *Acanthopagrus latus*", *Aquaculture*, 2020

Publication

<1 %

42

Venero, J.A.. "Variable feed allowance with constant protein input for the pacific white

<1 %

shrimp *Litopenaeus vannamei* reared under semi-intensive conditions in tanks and ponds", *Aquaculture*, 20070914

Publication

43

cicese.repositorioinstitucional.mx

Internet Source

<1 %

44

facultystaff.urmia.ac.ir

Internet Source

<1 %

45

link.springer.com

Internet Source

<1 %

46

www.scielo.br

Internet Source

<1 %

47

www.uanl.mx

Internet Source

<1 %

48

Dacley Hertes NEU, Wilson Rogério BOSCOLO, Fernanda Losi Alves d ALMEIDA, Micheli ZAMINHAN HASSEMER et al. "Growth performance, hematology, and muscle growth in isoleucine fed nile tilapia", *Boletim do Instituto de Pesca*, 2017

Publication

<1 %

49

Sudhansu Shekhar Mahanand, Sanjib Moulick, P. Srinivasa Rao. "Optimum formulation of feed for rohu, *Labeo rohita* (Hamilton), with biofloc as a component", *Aquaculture International*, 2012

Publication

<1 %

50

www.cabi.org

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On